

The Riemann–Roch theorem

Armand Borel and Jean-Pierre Serre

1958

Translator’s note

This page is a translation into English of the following:

Borel, A. and Serre, J.-P. “Le théorème de Riemann–Roch.” *Bulletin de la Société Mathématique de France* **86** (1958), 97–136. DOI: [10.24033/bsmf.1500](https://doi.org/10.24033/bsmf.1500).

The translator (Tim Hosgood) takes full responsibility for any errors introduced, and claims no rights to any of the mathematical content herein.

Version: [94f6dce](#)

[Translator] In the original article, subsections were numbered e.g. 5.a, 5.b, ...; here they are numbered 5.1, 5.2, ...

Contents

Introduction	2
1 Supplementary results concerning sheaves	2
2 Proper maps of quasi-projective varieties	4
3 Image of a sheaf under a proper map	6
4 The group $K(X)$ of classes of sheaves on a variety X	9
5 Operations on $K(X)$	12
5.1 Ring structure on $K(X)$	12
5.2 The exterior power operation	13
5.3 The operation $f^!$	14
5.4 The operation $f_!$	14
6 Chern classes	15
7 Statement of the Riemann–Roch theorem; First simplifications	16
8 Exactness and homotopy properties of $K(X)$	18
9 Proof of the Riemann–Roch theorem for $f: X \times \mathbf{P} \rightarrow X$	21

10 General remarks on the injection of a subvariety	22
10.1 Notation	22
10.2 Local parameters; Normal bundle	23
10.3 Local resolution of \mathcal{O}_Y over \mathcal{O}_X	23
10.4 —	24
10.5 Particular case of the divisor	26
11 Proof of R–R in the particular case of an injection	26
12 Blowing up along a subvariety	28
12.1 Notation	28
12.2 Definition of X' by local charts	28
12.3 Projective embedding of X'	29
13 Finishing the proof of R–R	30
14 Proof of parts (a) and (b) of Lemma 19	32
15 Proof of part (c) of Lemma 19	32
16 Proof of part (d) of Lemma 19	34
17 An application of R–R	36
Bibliography	37

Introduction

What follows constitutes the notes from a seminar that took place in Princeton in the autumn of 1957 on the works of Grothendieck; the new results that are included are due to Grothendieck; our contribution is solely of an editorial nature.

The “Riemann–Roch theorem” of which we speak here holds true for (non-singular) algebraic varieties over a field of arbitrary characteristic; in the classical case, where the base field is \mathbb{C} , this theorem encapsulates, as a particular example, the result proven a few years ago by Hirzebruch [10].

The full statement and proof of the Riemann–Roch theorem can be found in sections 7 to 16, with the last section being devoted to an application. Sections 1 to 6 contain some preliminaries on coherent algebraic sheaves [12]. The terminology that we follow is the same as in [12], up to one difference: to conform with a custom which is becoming more and more widespread, we use the word “morphism” instead of “regular map.”

1 Supplementary results concerning sheaves

(All the varieties considered below are algebraic varieties over an algebraically closed field k of arbitrary characteristic. Unless otherwise mentioned, all the sheaves considered are coherent algebraic sheaves.)

Proposition 1. *Let U be an open subset of a variety V , and let \mathcal{F} be a coherent sheaf on V and \mathcal{G} a coherent subsheaf of $\mathcal{F}|_U$ (the restriction of \mathcal{F} to U). Then there exists a coherent sheaf $\mathcal{G}' \subset \mathcal{F}$ such that $\mathcal{G}'|_U = \mathcal{G}$.*

(In fact, the proof will show that there exists a *largest* such sheaf having this property.)

Proof. For every open subset $W \subset V$, we define \mathcal{G}'_W as the set of sections of \mathcal{F} over W that belong to \mathcal{G} over $U \cap W$. Everything reduces to showing that the sheaf \mathcal{G}' associated to this presheaf is coherent. Since this is a local question, we can suppose that V is an affine variety. Let A be its coordinate ring. There exist elements $f_i \in A$ such that $U = \bigcup V_{f_i}$, where $V_{f_i} = U_i$ denotes the set of points of V where $f_i \neq 0$. If, in the definition of \mathcal{G}' , we replace the open subset U by the open subset U_i , then we obtain a sheaf $\mathcal{G}'_i \subset \mathcal{F}$, and it is clear that $\mathcal{G}' = \bigcap \mathcal{G}'_i$. By known results on coherent sheaves [12], it suffices to show that the \mathcal{G}'_i are coherent. We can thus restrict to considering the case where V is affine, and where $U = V_f$, for some $f \in A$. In this case, the sheaf \mathcal{F} is defined by an A -module M , and the subsheaf \mathcal{G} of $\mathcal{F}|_U$ is defined by a submodule N of $M_f = M \otimes_A A_f$. Let N' be the inverse image of N in M under the canonical map $M \rightarrow M_f$. The module N' then corresponds to a coherent subsheaf of \mathcal{F} , and we can immediately verify (by taking the $V_{f'}$ to be the W , for example) that this sheaf is exactly \mathcal{G}' , which finishes the proof. \square

Lemma 1. *Let U be an open subset of an affine variety V , and let \mathcal{F} be a (coherent) sheaf on U . Then \mathcal{F} is generated by its sections (over U).*

Proof. Let $x \in U$, and let f be a regular function on V , zero on $V \setminus U$ and non-zero at x . We have $V_f \subset U \subset V$. Since V_f is affine, we know [12] that \mathcal{F}_x is generated by its sections over V_f , and it thus suffices to prove that these sections can be extended to U , after multiplying by a suitable power of f . This follows from the following more general lemma: \square

Lemma 2. *Let X be a variety, f a regular function on X , \mathcal{F} a sheaf on X , and s a section of \mathcal{F} over $U = X_f$. Then there exists an integer $n > 0$ such that $f^n s$ can be extended to a section of \mathcal{F} over X .*

Proof. We can cover X by finitely many affine opens X_i . By applying [12, Lemma 1, p. 247], or by arguing directly (as in Proposition 1), we see that there exists an integer n and sections s_i of \mathcal{F} over the X_i that extend $f^n s$ over $X_i \cap U$. Since the $s_i - s_j$ are zero on $X_i \cap X_j \cap U$, there exists an integer m such that $f^m (s_i - s_j) = 0$ on $X_i \cap X_j$ ([12], or arguing directly), and m can be chosen independent of the pair (i, j) . The $f^m s_i$ then define a section s' of \mathcal{F} over X that indeed extend $f^{n+m} s$. \square

| p. 99

Proposition 2. *If U is an open subset of a variety V , then every sheaf \mathcal{F} over U can be extended to V .*

Proof. We show that, if $U \neq V$, we can extend \mathcal{F} to an open subset $U' \supset U$, with $U' \neq U$; from that fact that every (strictly) increasing chain of open subsets stabilises, this will imply the proposition. Let $x \in V \setminus U$, and let W be an affine open that contains x ; let

$U' = W \cup U$. We are thus led to extending the sheaf $\mathcal{F}|_{W \cap U}$ to W , or, in other words, we can restrict to proving the proposition in the specific case where V is affine. In this case, [Lemma 1](#) shows that \mathcal{F} is generated by its sections, i.e. it is of the form \mathcal{L}/\mathcal{R} , where \mathcal{L} is the direct sum of the sheaves \mathcal{O}_U . The sheaf \mathcal{L} can be extended in the obvious way to V , and, by [Proposition 1](#), there exists a subsheaf \mathcal{R}' of \mathcal{L} on V whose restriction to U is \mathcal{R} . The sheaf \mathcal{L}/\mathcal{R}' is then the desired extension. \square

Remark. [Proposition 1](#) and [Proposition 2](#) correspond to the geometric fact that the closure of any algebraic subvariety of U is an algebraic subvariety of V . These propositions do not extend *as is* to the *analytic* case. The most we can hope for (by results of Rothstein) is that they still hold true if we make certain restrictions on the dimensions of $V \setminus U$ and the varieties appearing in the local primary decomposition of the sheaf \mathcal{F} .

2 Proper maps of quasi-projective varieties

A variety X is said to be *quasi-projective* if it is isomorphic to a locally closed subvariety of a projective space. It is said to be *projective* if it is isomorphic to a closed subvariety of a projective space. *From here on in, all the varieties considered are assumed to be quasi-projective.*

Lemma 3. *Let \mathbf{P} be a projective space, U an arbitrary variety, and G a closed subset of $\mathbf{P} \times U$. Then the projection of G in U is closed.*

This is a translation into geometric language of the well-known fact that a projective space is a “complete” variety, in the sense of Weil. We briefly recall the principal of the proof:

Proof. Since the question is local with respect to U , we can assume that U is affine, and even that U is an affine open of the space k^n . We can also assume that G is irreducible. So we choose projective coordinates x_i in \mathbf{P} such that G meets the set $\mathbf{P}_0 \times U$ of points where $x_0 \neq 0$. If A denotes the coordinate ring of U , then the coordinate ring of the affine variety $\mathbf{P}_0 \times U$ is $A[x_i/x_0] = B_0$; the set G defines (and is defined by) a prime ideal \mathfrak{p} of B_0 . If \mathfrak{p}' denotes $A \cap \mathfrak{p}$, then the prime ideal \mathfrak{p}' corresponds to the closure of the projection G' of G in U . A point in this closure is thus a homomorphism $f: A \rightarrow k$ (where k denotes the base field) that is zero on \mathfrak{p}' ; this point is the image of a point in G that lies in $\mathbf{P}_0 \times U$ if and only if f can be extended to a homomorphism $g: B_0 \rightarrow k$ that is zero on \mathfrak{p} . So let L be the field of functions of G ; the field L contains A/\mathfrak{p}' as a subring. By the theorem of extension of specialisations, there exists a valuation v of L , with values in k , that extends f . Let Φ be the place associated to this valuation. If $v(x_i/x_0) \geq 0$ for all i , then the place Φ is finite over the x_i/x_0 , and thus induces, on $B_0/\mathfrak{p} \subset L$, a homomorphism g that extends f . If $v(x_i/x_0) < 0$ for some i , then we replace x_0 by the x_i that gives the smallest possible value of $v(x_i/x_0)$, and we are then back in the previous case. \square

| p. 100

If $f: X \rightarrow Y$ is a morphism, then we write G_f to denote its graph. It is trivial that G_f is closed in $X \times Y$.

Lemma 4. *Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be morphisms, where X and Y are subvarieties of projective spaces \mathbf{P} and \mathbf{P}' (respectively). Suppose that G_f is closed in $\mathbf{P} \times Y$, and that G_g is closed in $\mathbf{P}' \times Z$. Then G_{gf} is closed in $\mathbf{P} \times Z$.*

Proof. We have $G_f \subset \mathbf{P} \times Y = \mathbf{P} \times G_g \subset \mathbf{P} \times \mathbf{P}' \times Z$, and since each one is closed in the next, we see that G_f can be identified with a closed subset of $\mathbf{P} \times \mathbf{P}' \times Z$. Since G_{gf} is exactly the projection of G_f to the factor $\mathbf{P} \times Z$, the lemma follows from [Lemma 3](#). \square

Lemma 5. *Let $f: X \rightarrow Y$ be a morphism, and let $X \subset \mathbf{P}$ and $X \subset \mathbf{P}'$ be embeddings of X into projective spaces. If G_f is closed in $\mathbf{P} \times Y$, then it is also closed in $\mathbf{P}' \times Y$.*

Proof. We apply [Lemma 4](#) to the morphisms $X \xrightarrow{i} X \xrightarrow{f} Y$, where i denotes the identity morphism. Everything then reduces to showing that the graph G_i of i in $\mathbf{P} \times X$ is closed, which follows from the fact that it is given by the intersection of $\mathbf{P} \times X$ with the diagonal of $\mathbf{P} \times \mathbf{P}$. \square

[Lemma 5](#) justifies the following definition:

Definition. A map $f: X \rightarrow Y$ is said to be *proper* if it is a morphism and if its graph G_f is closed in $\mathbf{P} \times Y$, where \mathbf{P} is a projective space containing X .

We can give a definition of proper maps that is analogous to the definition of complete varieties:

Proposition 3. *For a morphism $f: X \rightarrow Y$ to be proper, it is necessary and sufficient, for every variety Z and every closed subset T of $X \times Z$, for the image of T in $Y \times Z$ to be closed.*

Proof. Let \mathbf{P} be a projective space inside which X can be embedded; since G_f is closed in $\mathbf{P} \times Y$, the product $G_f \times Z$ is closed in $\mathbf{P} \times Y \times Z$, and so T can be embedded as a closed subset into $\mathbf{P} \times Y \times Z$. Applying [Lemma 3](#), we see that the projection of T to $Y \times Z$ (which is exactly $(f \times 1)(T)$) is closed. Conversely, suppose that this property holds true, and apply it to $Z = \mathbf{P}$, with the set T being the diagonal of $X \times X$, embedded into $X \times \mathbf{P}$. The image of T in $Y \times Z = Y \times \mathbf{P}$ is then exactly G_f , which is indeed closed. \square

| p. 101

Proposition 4.

—

- i. *The identity morphism $i: X \rightarrow X$ is proper.*
- ii. *The composition of two proper maps is proper.*
- iii. *The direct product of two proper maps is proper.*
- iv. *The image of a closed subset by a proper map is a closed subset.*
- v. *An injection $Y \rightarrow X$ is proper if and only if Y is closed in X .*
- vi. *Every morphism from a projective variety is proper.*
- vii. *A projection $Y \times Z \rightarrow Y$ is proper if and only if Z is projective (assuming the variety Y to be non-empty).*

Proof. We indicate, as an example, how to prove (vii) (since the other claims are even easier to prove). If Z is projective, then we apply the criteria of [Proposition 3](#); so let Z' be an arbitrary variety, and T a closed subset of $Y \times Z \times Z'$; we need to show that the projection of T in $Y \times Z'$ is closed, which follows from [Lemma 3](#). Conversely, if $Y \times Z \rightarrow Y$ is proper, then the composition $Z \rightarrow Y \times Z \rightarrow Y$ is proper. Since the image of this map is a point, we immediately deduce that Z is projective (by returning to the definition). \square

(*[Trans.]* The following corollary is labelled “Corollary 5” in the original, but this seems to be a numbering error).

Corollary. *For a morphism $f: X \rightarrow Y$ to be proper, it is necessary and sufficient for it to factor as $X \rightarrow \mathbf{P} \times Y \rightarrow Y$, where $X \rightarrow \mathbf{P} \times Y$ is an injection into a closed subvariety, and $\mathbf{P} \times Y \rightarrow Y$ is the projection onto the second factor (where \mathbf{P} denotes some projective space).*

Proof. By the definition of a proper map, this condition is necessary (if we take \mathbf{P} to be a projective space into which we can embed X). It is sufficient by (ii), (v), and (vii). \square

Proposition 5. *Suppose that the base field k is the field of complex numbers. For a morphism $f: X \rightarrow Y$ to be proper (in the above sense), it is necessary and sufficient for it to be proper (in the topological sense) when we endow X and Y with the “usual” topology.*

Proof. Suppose that f is proper in the algebraic sense, and let K be a compact subset of Y (for the usual topology). Suppose that X is embedded into some projective space \mathbf{P} . Since \mathbf{P} is compact, we know that $f^{-1}(K) = G_f \cap (\mathbf{P} \times K)$ is compact, which shows that f is proper in the topological sense. Conversely, assume that this condition is satisfied, and aim to prove that the condition of [Proposition 3](#) is satisfied: the image of T in $Y \times Z$ is closed for the usual topology, and thus also for the Zariski topology [[13](#), [Proposition 7](#), p. 12]. \square

| p. 102

Remark. The notion of a proper map can be extended to “abstract” (that is, non-quasi-projective) varieties: it suffices to take the criteria of [Proposition 3](#) as a definition [[4](#)]. [Proposition 4](#) and [Proposition 5](#) still hold true (if we replace “projective” with “complete” in item (vii) of [Proposition 4](#)). The proofs are essentially the same: instead of using embeddings into projective spaces, we use the fact that every variety is the image of a quasi-projective variety under a proper map (Chow’s lemma [[4](#),[14](#)]).

3 Image of a sheaf under a proper map

Let $f: X \rightarrow Y$ be a morphism, with Y a variety, and let \mathcal{F} be a (coherent algebraic, as always) sheaf on X . We define, by the classical procedure of Leray, sheaves $R^q f(\mathcal{F})$ on Y by setting

$$R^q f(\mathcal{F})_U = H^q(f^{-1}(U), \mathcal{F}) \quad \text{for every open subset } U \text{ of } Y.$$

For $q = 0$, we have the sheaf associated to the presheaf given by the $H^0(f^{-1}(U), \mathcal{F})$; this is the *direct image* of the sheaf \mathcal{F} . We can show [[8](#)] that the $R^q f$ are the *derived functors* of

the functor $\mathcal{F} \rightarrow R^0 f(\mathcal{F})$ (where \mathcal{F} runs over the category of all sheaves on X , coherent or not).

Examples.

1. If $X \rightarrow Y$ is an injection of a closed subvariety, then the sheaf $R^0 f(\mathcal{F})$ is exactly the sheaf \mathcal{F} extended by 0 outside of X , and the sheaves $R^q f(\mathcal{F})$, for $q \geq 1$, are zero (let U be an affine open; then $f^{-1}(U) = U \cap X$ is affine, whence $R^q f(\mathcal{F}) = 0$).
2. Let Y be a point. A sheaf on a point is simply a group (or a k -vector space, if we are talking about algebraic sheaves). The $R^q f(\mathcal{F})$ are then simply the cohomology groups $H^q(X, \mathcal{F})$; we note that these are not necessarily vector spaces of *finite dimension* (or, in other words, not necessarily *coherent* sheaves on Y).
3. Suppose that $f: X \rightarrow Y$ defines a birational isomorphism between the varieties X and Y (assumed to be projective and non-singular). Take \mathcal{F} to be the sheaf \mathcal{O}_X of local rings of X ; we immediately see that $R^0 f(\mathcal{O}_X) = \mathcal{O}_Y$. Is it true that $R^q f(\mathcal{O}_X) = 0$ for $q \geq 1$? We can at least verify this for “blow-ups,” and it would be interesting to know the answer in the general case.

We note that Leray’s theory can be translated without any changes (see [8]); there is a spectral sequence abutting $H^*(X, \mathcal{F})$, and with $E_2^{p,q} = H^p(Y, R^q f(\mathcal{F}))$. If we apply, for example, this spectral sequence to [Example 3](#) above, then we see that $R^q f(\mathcal{O}_X) = 0$ for $q \geq 1$ implies that $H^*(X, \mathcal{O}_X) = H^*(Y, \mathcal{O}_Y)$.

| p. 103

We have seen ([Example 2](#)) that the $R^q f(\mathcal{F})$ are not, in general, coherent sheaves on Y . However:

Theorem 1. *If $f: X \rightarrow Y$ is proper, then the $R^q f(\mathcal{F})$, for $q > 0$, are coherent sheaves on Y , for any coherent sheaf \mathcal{F} on X .*

Let \mathbf{P} be a projective space into which we can embed X , and let G_f be the graph of f in $\mathbf{P} \times Y$. By definition of what it means to be a proper map, G_f is closed in $\mathbf{P} \times Y$. Let \mathcal{F}' be the sheaf on $\mathbf{P} \times Y$ obtained by extending \mathcal{F} by 0 outside of $G_f = X$ (see [Example 2](#)); if π denotes the projection from $\mathbf{P} \times Y$ to Y , then we immediately see that $R^q \pi(\mathcal{F}') = R^q f(\mathcal{F})$. We can thus restrict to proving [Theorem 1](#) for $\pi: \mathbf{P} \times Y \rightarrow Y$. Furthermore, since the question is local with respect to Y , we can assume that Y is an affine variety.

On \mathbf{P} we have a “standard” bundle L of dimension 1, whose sections are the linear forms (see [12, Chap. III, §2]); this fibre defines a fibre on $X = \mathbf{P} \times Y$ that we denote also by L . The sheaf associated to L^n on $\mathbf{P} \times Y$ will be denoted $\mathcal{O}_X(n)$. We then have:

Lemma 6. *Every coherent algebraic sheaf \mathcal{F} on $X = \mathbf{P} \times Y$, with Y affine, is isomorphic to a quotient of some direct sum of sheaves of the form $\mathcal{O}_X(n)$.*

Proof. When Y consists of a single point, this is [12, Théorème 1, p. 247]. We are going to reduce to this particular case: let $Y \subset \mathbf{P}'$ be an embedding of Y into a projective space, and let \bar{Y} be the closure of Y . By [Proposition 2](#), the sheaf \mathcal{F} can be extended to a sheaf $\bar{\mathcal{F}}$ on $\mathbf{P} \times \bar{Y}$. The embedding of \bar{Y} into \mathbf{P}' defines, on \bar{Y} (and thus also on $\mathbf{P} \times \bar{Y}$), a bundle L' of dimension 1. The product bundle LL' corresponds to the well known embedding of

$\mathbf{P} \times \mathbf{P}'$ into a projective space P'' (the ‘‘Segre’’ embedding, given by the products $x_i y_j$ of the homogeneous coordinates of the two projective spaces). Then applying the result from [12] cited above to \mathcal{F} and \mathbf{P}'' , we see that \mathcal{F} is a quotient of a direct sum of sheaves of the form $\mathcal{O}_{\mathbf{P} \times \overline{Y}}(L^n L'^m)$; by restricting to $\mathbf{P} \times Y$, and taking into account the fact that \mathbf{P}' is *trivial* on Y , we indeed obtain the desired result. \square

[Of course, we could also give a direct proof, copied from the one in [12].]

Lemma 7. *The $R^q \pi(\mathcal{O}_X(n))$ are coherent sheaves on Y .*

Proof. We explicitly calculate the sheaves $R^q \pi(\mathcal{O}_X(n))$. If U is an affine open subset of Y , then | p. 104

$$R^q \pi(\mathcal{O}_X(n))_U = H^q(\mathbf{P} \times U, \mathcal{O}_X(n)).$$

If $\mathcal{U} = \{U_i\}$ is an affine cover of \mathbf{P} , then the $U_i \times U$ form an affine cover \mathcal{U}' of $\mathbf{P} \times U$; taking into account the fact that $\mathcal{O}_X(n)$ ‘‘comes from’’ \mathbf{P} , we see that the complex $C(\mathcal{U}', \mathcal{O}_X(n))$ is isomorphic to the tensor product $C(\mathcal{U}', \mathcal{O}_{\mathbf{P}}(n)) \otimes_k H^0(U, \mathcal{O}_U)$. The universal coefficient formula then shows that

$$H^q(\mathbf{P} \times U, \mathcal{O}_X(n)) = H^q(\mathbf{P}, \mathcal{O}_{\mathbf{P}}(n)) \otimes_k H^0(U, \mathcal{O}_U).$$

This latter equality implies that $R^q \pi(\mathcal{O}_X(n))$ is isomorphic to the sheaf $\mathcal{O}_Y \otimes_k V^q$, where $V^q = H^q(\mathbf{P}, \mathcal{O}_{\mathbf{P}}(n))$. Since V^q is a vector space of finite dimension over k , it is indeed a coherent sheaf on Y , which proves the lemma. \square

[The above proof applies more generally to any projection $\pi: Y \times Z \rightarrow Y$ with Z projective, whenever the sheaf \mathcal{F} is of the form $\mathcal{G} \otimes \mathcal{H}$, with \mathcal{G} coherent on Y and \mathcal{H} coherent on Z . We then find that $R^q \pi(\mathcal{F}) = \mathcal{G} \otimes H^q(Z, \mathcal{H})$. We could consider, even more generally, the case of a product map $Y \times Z \rightarrow Y' \times Z' \dots$]

Proof. (of Theorem 1). We can now prove Theorem 1 for an arbitrary sheaf \mathcal{F} on $X = \mathbf{P} \times Y$. We argue by decreasing induction on the integer q . If $q > \dim X$, then it is clear that $R^q \pi(\mathcal{F}) = 0$. So suppose that the theorem is proven for $q + 1$. By Lemma 6, there exists an exact sequence $0 \rightarrow \mathcal{R} \rightarrow \mathcal{L} \rightarrow \mathcal{F} \rightarrow 0$, where \mathcal{L} is isomorphic to a direct sum of sheaves of the form $\mathcal{O}_X(n)$. The exact sequence of cohomology (or the exact sequence of derived functors) shows that we have an exact sequence

$$R^q \pi(\mathcal{R}) \rightarrow R^q \pi(\mathcal{L}) \rightarrow R^q \pi(\mathcal{F}) \rightarrow R^{q+1} \pi(\mathcal{R}) \rightarrow R^{q+1} \pi(\mathcal{L}).$$

Given the induction hypothesis and Lemma 7, the sheaves $R^q \pi(\mathcal{L})$, $R^{q+1} \pi(\mathcal{R})$, and $R^{q+1} \pi(\mathcal{L})$ are coherent. It follows that $R^q \pi(\mathcal{F})$ admits a subsheaf of finite type, with the quotient being coherent. An immediate argument then shows that $R^q \pi(\mathcal{F})$ is of finite type. This result, having been proven for any coherent sheaf, also holds for \mathcal{R} . The image of $R^q \pi(\mathcal{R})$ in $R^q \pi(\mathcal{L})$ is then a coherent sheaf ([12]), and $R^q \pi(\mathcal{F})$ is an extension of two coherent sheaves, and is thus coherent (*id.*). \square

Remarks.

—

1. **Theorem 1** holds true even if we don't suppose that X is quasi-projective (we can restrict to this case by using Chow's lemma and the "devissage" of coherent sheaves, see [7]).
2. Grauert and Remmert have proven the analytic analogue of **Theorem 1** for the projection $\pi: \mathbf{P} \times Y \rightarrow Y$. Needless to say that the proof is more difficult!

4 The group $K(X)$ of classes of sheaves on a variety X

Let X be an algebraic variety, and let $F(X)$ be the free abelian group generated by the set \mathcal{C} of (coherent algebraic, as always) sheaves on X . An element of $F(X)$ is thus a formal linear combination

$$x = \sum n_i \mathcal{F}_i \quad \text{where } n_i \in \mathbb{Z} \text{ and } \mathcal{F}_i \in \mathcal{C}.$$

We agree, of course, to identify isomorphic sheaves [if not, then $F(X)$ would not even be a "set!"].

Let

$$(E) \quad 0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$$

be an exact sequence of sheaves. To this exact sequence we associate the element $Q(E) = \mathcal{F} - \mathcal{F}' - \mathcal{F}''$ of $F(X)$.

Definition. We define the *group of classes of sheaves on X* to be the quotient of $F(X)$ by the subgroup generated by the $Q(E)$, where E runs over all short exact sequences.

This group will be denoted by $K(X)$ in what follows. If \mathcal{F} is a sheaf on X , then its canonical image in $K(X)$ will be denoted $\gamma_X(\mathcal{F})$, or $\gamma(\mathcal{F})$, or simply \mathcal{F} , depending on the risk of confusion. The $\gamma(\mathcal{F})$ generate $K(X)$, and the map $\mathcal{F} \rightarrow \gamma(\mathcal{F})$ is "additive"; in other words, if we have the exact sequence (E), then $\gamma(\mathcal{F}) = \gamma(\mathcal{F}') + \gamma(\mathcal{F}'')$. Conversely, by the very definition of $K(X)$, every additive map from the set of sheaves into an abelian group G can be written in the form $\mathcal{F} \mapsto \pi(\gamma(\mathcal{F}))$, where $\pi: K(X) \rightarrow G$ is a uniquely determined homomorphism.

We can apply the above to construction to many other situations, apart from that of sheaves. We will need, in particular, to apply it to the case of *vector bundles* on X . So let \mathcal{V} be the set of vector bundles; we define $F_1(X)$ to be the free group generated by \mathcal{V} , and $K_1(X)$ to be the quotient of $F_1(X)$ by the subgroup generated by the $Q_1(E) = \mathcal{F} - \mathcal{F}' - \mathcal{F}''$, where (E) now denotes a short exact sequence of vector bundles. If X is connected (which we will assume), then we know that we can identify vector bundles with locally free sheaves on X ; we thus have $\mathcal{V} \subset \mathcal{C}$, and the injection $\mathcal{V} \rightarrow \mathcal{C}$ defines a canonical homomorphism $\varepsilon: K_1(X) \rightarrow K(X)$.

Theorem 2. *Suppose that X is an irreducible non-singular quasi-projective variety. Then the homomorphism $\varepsilon: K_1(X) \rightarrow K(X)$ defined above is a bijection.*

We will need a certain number of auxiliary results on the relation between \mathcal{V} and \mathcal{C} :

Lemma 8. *Let $0 \rightarrow \mathcal{L} \rightarrow \mathcal{L}' \rightarrow \mathcal{L} \rightarrow 0$ be an exact sequence such that $\mathcal{L}', \mathcal{L} \in \mathcal{V}$. Then $\mathcal{L} \in \mathcal{V}$.*

Proof. If $P \in X$, then the local module \mathcal{L}_P is free over \mathcal{O}_P , and so a direct factor of \mathcal{L}'_P , which proves that \mathcal{L}_P is a projective \mathcal{O}_P -module, and thus free, since \mathcal{O}_P is a local ring. But a coherent algebraic sheaf such that all its stalks are free is itself locally free (see [12, lines 10–11 from the bottom]). \square

Lemma 9. *Let $n = \dim X$, and let $0 \rightarrow \mathcal{L} \rightarrow \mathcal{L}_p \rightarrow \dots \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0$ be an exact sequence, with $\mathcal{L}_i \in \mathcal{V}$. If $p \geq n - 1$, then $\mathcal{L} \in \mathcal{V}$.*

Proof. This is again a local question. So let $P \in X$; the fact that the local ring \mathcal{O}_P is a regular local ring of dimension n means that we can apply the syzygy theorem, and show that \mathcal{L}_P is \mathcal{O}_P -free, whence the desired result (noting that the hypothesis that X is non-singular is used in an essential way). \square

Lemma 10. *Every $\mathcal{F} \in \mathcal{C}$ is a quotient of some $\mathcal{L} \in \mathcal{V}$.*

Proof. Let $X \subset \mathbf{P}$ be a projective embedding of X , and let \bar{X} be its closure in \mathbf{P} . By [Proposition 2](#), \mathcal{F} can be extended to a sheaf \mathcal{F}' on \bar{X} . By [12, Théorème 1, p. 247] (and see also [Lemma 6](#)), the sheaf \mathcal{F}' is the quotient of a direct sum of sheaves of the form $\mathcal{O}_{\bar{X}}(n)$, and thus a locally free sheaf on \bar{X} . By restriction to X , we obtain the desired result. \square

Corollary. *For every $\mathcal{F} \in \mathcal{C}$, there exists an exact sequence*

$$0 \rightarrow \mathcal{L}_n \rightarrow \mathcal{L}_{n-1} \rightarrow \dots \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0$$

with $\mathcal{L}_i \in \mathcal{V}$.

Proof. This is a consequence of [Lemma 9](#) and [Lemma 10](#). We can state this corollary in a different way by saying that there exists a “complex” \mathcal{L} in \mathcal{V} that is acyclic in degrees ≥ 1 , and such that $H_0(\mathcal{L}) = \mathcal{F}$. \square

Proof. (of [Theorem 2](#)). We now continue to the proof of [Theorem 2](#). If $\mathcal{F} \in \mathcal{C}$, we take an acyclic complex \mathcal{L} in \mathcal{V} such that $H_0(\mathcal{L}) = \mathcal{F}$. We set

$$\gamma_1(\mathcal{L}) = \sum (-1)^p \gamma_1(\mathcal{L}_p);$$

this is an element of $K_1(X)$; we define $\gamma(\mathcal{L}) \in K(X)$ analogously. Suppose that the following two lemmas have been proven:

Lemma 11. $\gamma_1(\mathcal{L})$ depends only on \mathcal{F} .

Lemma 12. $\gamma_1(\mathcal{L})$ is an additive function of \mathcal{F} .

Then, setting $\eta(\mathcal{F}) = \gamma_1(\mathcal{L})$, we obtain a homomorphism $\eta: K(X) \rightarrow K_1(X)$. By the definition of \mathcal{L} , we have that

$$\eta(\gamma_1(\mathcal{L})) = \gamma(\mathcal{L}) = \gamma(\mathcal{F}),$$

whence $\varepsilon \circ \eta = 1$. The proof that $\eta \circ \varepsilon = 1$ is even more trivial. So everything relies only on proving [Lemma 11](#) and [Lemma 12](#). \square | p. 107

Proof. (of [Lemma 11](#)). We first start by stating a corollary of [Lemma 10](#):

Lemma 13. *Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathcal{C}$, and let $u: \mathcal{A} \rightarrow \mathcal{B}$ and $v: \mathcal{C} \rightarrow \mathcal{B}$, with u and v surjective. Then there exist $\mathcal{L} \in \mathcal{V}$, and $u': \mathcal{L} \rightarrow \mathcal{C}$ and $v': \mathcal{L} \rightarrow \mathcal{A}$, such that $v \circ u' = u \circ v'$, with u' and v' surjective:*

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{v'} & \mathcal{A} \\ u' \downarrow & & \downarrow u \\ \mathcal{C} & \xrightarrow{v \circ u'} & \mathcal{B} \end{array}$$

Proof. Let $(\mathcal{A}, \mathcal{C})$ be the subsheaf of $\mathcal{A} \times \mathcal{C}$ given by the elements having the same image in \mathcal{B} . Since u and v are surjective, the canonical projections $(\mathcal{A}, \mathcal{C}) \rightarrow \mathcal{A}$ and $(\mathcal{A}, \mathcal{C}) \rightarrow \mathcal{C}$ are surjective. By applying [Lemma 10](#) to $(\mathcal{A}, \mathcal{C})$, we obtain the desired result. \square

Now let \mathcal{L} and \mathcal{L}' be two resolutions of \mathcal{F} , which we intend to compare. We will show that there exists a third resolution \mathcal{L}'' of \mathcal{F} , along with surjective homomorphisms $\mathcal{L}'' \rightarrow \mathcal{L}$ and $\mathcal{L}'' \rightarrow \mathcal{L}'$ that induce the identity on H^0 ; everything will then rely only on proving that $\gamma_1(\mathcal{L}'') = \gamma_1(\mathcal{L})$, for example. But, if we denote by \mathcal{L}_1 the kernel of $\mathcal{L}'' \rightarrow \mathcal{L}$, then $\mathcal{L}_1 \in \mathcal{V}$, by [Lemma 8](#), and the exact sequence of homology shows that $H^q(\mathcal{L}_1) = 0$ for all $q \geq 0$. We thus immediately deduce that $\gamma_1(\mathcal{L}_1) = 0$, and, since $\gamma_1(\mathcal{L}'') = \gamma_1(\mathcal{L}) + \gamma_1(\mathcal{L}_1)$, this gives the desired result. Everything thus relies only on proving the existence of the resolution \mathcal{L}'' , which we do dimension by dimension (where [Lemma 9](#) tells us that the procedure will eventually stop), using the following lemma:

Lemma 14. *Let*

$$0 \rightarrow \mathcal{L}'' \rightarrow \mathcal{L} \rightarrow \mathcal{B} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \mathcal{L}'' \rightarrow \mathcal{L}' \rightarrow \mathcal{B}' \rightarrow 0$$

be exact sequences (with $\mathcal{L}, \mathcal{L}' \in \mathcal{V}$), and let $\mathcal{B}'' \rightarrow \mathcal{B}$ and $\mathcal{B}'' \rightarrow \mathcal{B}'$ be surjective maps. We can then extend \mathcal{B}'' to an exact sequence

$$0 \rightarrow \mathcal{L}'' \rightarrow \mathcal{L}'' \rightarrow \mathcal{B}'' \rightarrow 0 \quad \text{with } \mathcal{L}'' \in \mathcal{V}$$

and find surjective homomorphisms

$$\mathcal{L}'' \rightarrow \mathcal{L}, \quad \mathcal{L}'' \rightarrow \mathcal{L}', \quad \mathcal{L}'' \rightarrow \mathcal{L}, \quad \mathcal{L}'' \rightarrow \mathcal{L}'$$

such that the following diagram commutes:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{L} & \longrightarrow & \mathcal{L} & \longrightarrow & \mathcal{B} & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & \mathcal{L}'' & \longrightarrow & \mathcal{L}'' & \longrightarrow & \mathcal{B}'' & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{L}' & \longrightarrow & \mathcal{L}' & \longrightarrow & \mathcal{B}' & \longrightarrow & 0 \end{array}$$

Proof. By applying [Lemma 13](#) to $\mathcal{L} \rightarrow \mathcal{B}$ and $\mathcal{B}'' \rightarrow \mathcal{B}$, we see that $\mathcal{L}_1 \rightarrow \mathcal{L}$ and $\mathcal{L}_1 \rightarrow \mathcal{B}''$ are surjective. By applying the same lemma to $\mathcal{L}_1 \rightarrow \mathcal{B}'' \rightarrow \mathcal{B}'$ and $\mathcal{L}' \rightarrow \mathcal{B}'$, we see that $\mathcal{L}_2 \rightarrow \mathcal{L}_1$ and $\mathcal{L}_2 \rightarrow \mathcal{L}'$ are surjective, making the diagram commute.

Now let $\mathcal{L}_3 \rightarrow \mathcal{L}$ and $\mathcal{L}'_3 \rightarrow \mathcal{L}'$ be surjective, and set

$$\mathcal{L}'' = \mathcal{L}_2 \oplus \mathcal{L}_3 \oplus \mathcal{L}'_3.$$

We define $\mathcal{L}'' \rightarrow \mathcal{B}''$ as being 0 on \mathcal{L}_3 and \mathcal{L}'_3 , and equal to $\mathcal{L}_2 \rightarrow \mathcal{L}_1 \rightarrow \mathcal{B}$ on \mathcal{L}_2 ; we define $\mathcal{L}'' \rightarrow \mathcal{L}$ as being equal to $\mathcal{L}_2 \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}$ on \mathcal{L}_2 , to 0 on \mathcal{L}'_3 , and to $\mathcal{L}_2 \rightarrow \mathcal{L} \rightarrow \mathcal{L}$ on \mathcal{L}_3 ; we define $\mathcal{L}'' \rightarrow \mathcal{L}'$ analogously. We then define \mathcal{F}'' as the kernel of $\mathcal{L}'' \rightarrow \mathcal{B}''$, and $\mathcal{F}'' \rightarrow \mathcal{L}$ and $\mathcal{F}'' \rightarrow \mathcal{L}'$ as restrictions of the maps from \mathcal{L}'' . The commutativity of the diagram is then immediate; furthermore, \mathcal{F}'' evidently contains \mathcal{L}_3 , which maps to \mathcal{L} ; *a fortiori*, $\mathcal{F}'' \rightarrow \mathcal{L}$ is surjective, and so too is $\mathcal{F}'' \rightarrow \mathcal{L}'$, which finished the proof of [Lemma 13](#). \square

This finishes the proof of [Lemma 11](#). \square

Proof. (of [Lemma 12](#)). Let $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ be an exact sequence. We are going to show that there exists resolutions \mathcal{L}' , \mathcal{L} , and \mathcal{L}'' of these sheaves, such that we also have an exact sequence $0 \rightarrow \mathcal{L}' \rightarrow \mathcal{L} \rightarrow \mathcal{L}'' \rightarrow 0$. The additivity of $\gamma_1(\mathcal{L})$ will then be evident. To construct these resolutions, we proceed, again, dimension by dimension. Everything relies on proving that, given an exact sequence as above, we can extend it to a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{F}' & \longrightarrow & \mathcal{F} & \longrightarrow & \mathcal{F}'' & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & \mathcal{L}' & \longrightarrow & \mathcal{L} & \longrightarrow & \mathcal{L}'' & \longrightarrow & 0 \end{array}$$

with $\mathcal{L}', \mathcal{L}, \mathcal{L}'' \in \mathcal{V}$, and where the $\mathcal{L} \rightarrow \mathcal{F}$ are surjective.

For this, we first take some surjective $\mathcal{L}'' \rightarrow \mathcal{F}''$, and apply [Lemma 13](#) to $\mathcal{F} \rightarrow \mathcal{F}''$ and $\mathcal{L} \rightarrow \mathcal{F}''$. From this we obtain surjective $\mathcal{L}_1 \rightarrow \mathcal{F}$ and $\mathcal{L}_1 \rightarrow \mathcal{L}''$ that make the diagram commute. Next, we take some surjective $\mathcal{L}_2 \rightarrow \mathcal{F}'$, and set $\mathcal{L} = \mathcal{L}_2 \oplus \mathcal{L}_1$. We define $\mathcal{L} \rightarrow \mathcal{F}$ and $\mathcal{L} \rightarrow \mathcal{L}''$ in the evident way, and we take \mathcal{L}' to be the kernel of $\mathcal{L} \rightarrow \mathcal{L}''$. Then $\mathcal{L}_2 \subset \mathcal{L}'$, which shows that \mathcal{L}' maps to \mathcal{F}' , and all the desired conditions are satisfied. \square

This completes the proof of [Theorem 2](#).

Remark. The hypothesis that X is quasi-projective was used only in [Lemma 10](#), to prove that every coherent sheaf on X is the quotient of some locally free sheaf. We do not know if this lemma can be extended to “abstract” algebraic varieties.

5 Operations on $K(X)$

5.1 Ring structure on $K(X)$

Let \mathcal{F} and \mathcal{G} be (coherent) sheaves on X . The $\mathrm{Tor}_p(\mathcal{F}, \mathcal{G})$ ($p = 0, 1, \dots$) are coherent sheaves on X (of course, the Tor are taken over the structure sheaf of X). Since X is

non-singular, $\text{Tor}_p(\mathcal{F}, \mathcal{G}) = 0$ when $p > \dim X$, which means that the alternating sum $\chi(\mathcal{F}, \mathcal{G}) = \sum (-1)^p \text{Tor}_p(\mathcal{F}, \mathcal{G})$ is a well-defined element of $K(X)$. The exact sequence of Tor shows that $\chi(\mathcal{F}, \mathcal{G})$ is bilinear in \mathcal{F} and \mathcal{G} , and so extends to a bilinear map from $K(X) \times K(X)$ to $K(X)$, that we will denote by $(x, x') \mapsto x \cdot x'$.

Proposition 6. *The product defined above is commutative and associative.*

Proof. Commutativity is trivial, since each Tor is commutative. The associativity follows from the “associativity formula” of the Tor (see [3]): we define the “simultaneous Tor” $\text{Tor}(\mathcal{F}, \mathcal{G}, \mathcal{H})$, and two spectral sequences abutting to $\text{Tor}(\mathcal{F}, \mathcal{G}, \mathcal{H})$, and with E_2 pages $\text{Tor}(\mathcal{F}, \text{Tor}(\mathcal{G}, \mathcal{H}))$ and $\text{Tor}(\text{Tor}(\mathcal{F}, \mathcal{G}), \mathcal{H})$ (respectively); we use the fact that the Euler-Poincaré characteristics are invariant in a spectral sequence.

We can give a simpler proof by using [Theorem 2](#): note that, if \mathcal{F} and \mathcal{G} are locally free, then

$$\chi(\mathcal{F}, \mathcal{G}) = \gamma(\mathcal{F} \otimes \mathcal{G})$$

which means that associativity is evident when \mathcal{F}, \mathcal{G} , and \mathcal{H} are locally free. Since $K(X)$ is generated by the $\gamma(\mathcal{F})$, with \mathcal{F} locally free, this proves associativity. \square

[The above product thus corresponds to the tensor product of vector bundles.]

5.2 The exterior power operation

Let E be a vector bundle. The exterior powers $\wedge^p E$ are vector bundles, and their class in $K(X) = K_1(X)$ will be denoted by $\lambda^p(E)$. If we have an exact sequence

$$0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$$

then we define, by a well known procedure (Koszul’s thesis!) a filtration of $\wedge(E)$ with quotients $\wedge(E') \otimes \wedge(E'')$. From this we obtain the following formula:

$$\lambda^p(E) = \sum_{r+s=p} \lambda^r(E') \cdot \lambda^s(E'').$$

This formula can be understood as an additivity formula by introducing the formal series (in t)

$$\lambda_t(E) = \sum \lambda^p(E) t^p;$$

this is an element of $K(X)[[t]]$, starting with 1. The above formula implies that

| p. 110

$$\lambda_t(E) = \lambda_t(E') \lambda_t(E'').$$

The map $E \mapsto \lambda_t(E)$ can thus be extended to a homomorphism $x \mapsto \lambda_t$ from $K(X) = K_1(X)$ to the multiplicative group U of formal series

$$1 + a_1 t + \dots + a_n t^n + \dots \quad \text{with } a_i \in K(X).$$

By definition, $\lambda^p(x)$ is the coefficient of t^p in $\lambda_t(x)$.

In characteristic 0, Grothendieck has shown that, for every sheaf \mathcal{F} , $\lambda^p(\mathcal{F})$ is equal to the alternating sum of the “alternating Tor” of p copies of \mathcal{F} . In characteristic $\neq 0$, we do not know any analogous formula.

5.3 The operation $f^!$

Let $f: Y \rightarrow X$ be a morphism. If E is a vector bundle over X , then the bundle $f^{-1}(E)$ is a vector bundle over Y . This operation is additive, and thus extends to a homomorphism $f^!: K(X) \rightarrow K(Y)$. By arguing on the fibres, we immediately see that $f^!$ is a ring homomorphism, compatible with the operation λ^p , and such that $(fg)^! = g^!f^!$.

If \mathcal{F} is a coherent sheaf on X , then we can directly define $f^!(\mathcal{F})$ as the alternating sum of the $\text{Tor}_p^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{F})$; indeed, this expression is additive in \mathcal{F} (by the exact sequence of Tor), and reduces to $\mathcal{O}_Y \otimes \mathcal{F}$ when \mathcal{F} is locally free.

5.4 The operation $f_!$

Again, let $f: Y \rightarrow X$ be a morphism, that we now assume to be *proper*. If \mathcal{F} is coherent on Y , then we have seen (Theorem 1, §3) that the $R^q f(\mathcal{F})$ ($q = 0, 1, \dots$) are coherent sheaves on X , and their alternating sum is a well-defined element of $K(X)$. Since this alternating sum is additive in \mathcal{F} (by the exact sequence of cohomology), we thus obtain an additive homomorphism $f_!: K(Y) \rightarrow K(X)$. In the particular case where Y is a closed subvariety of X , and f is the canonical injection $Y \rightarrow X$, this operation reduces to extending by 0 outside of Y .

The map $f_!$ is not compatible with multiplication. We do, however, have the following formula:

$$f_!(y \cdot f^!(x)) = f_!(y) \cdot x \quad \text{for } x \in K(X) \text{ and } y \in K(Y).$$

It suffices to prove this formula when $y = \gamma_Y(\mathcal{F})$ and $x = \gamma_X(\mathcal{L})$, where \mathcal{F} (resp. \mathcal{L}) is a coherent sheaf on Y (resp. a locally free sheaf on X). In this case, we even have the more precise formula

$$R^q f(\mathcal{F} \otimes_{\mathcal{O}_Y} f^{-1}(\mathcal{L})) = R^q f(\mathcal{F}) \otimes_{\mathcal{O}_X} \mathcal{L} \quad (\star)$$

where we set

$$f^{-1}(\mathcal{L}) = \mathcal{L} \otimes_{\mathcal{O}_X} \mathcal{O}_Y.$$

To prove (\star) , we first note that

$$\mathcal{F} \otimes_{\mathcal{O}_Y} f^{-1}(\mathcal{L}) = \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}.$$

Returning to the definition of $R^q f$, we define a canonical homomorphism from the right-hand side of (\star) to the left-hand side; to show that it is an isomorphism we can argue locally. We can thus restrict to the case where $\mathcal{L} = \mathcal{O}_X$, and in this case our claim is trivial.

If we have two proper maps $Z \xrightarrow{g} Y \xrightarrow{f} X$, and if \mathcal{F} is a coherent sheaf on Z , then we can construct a spectral sequence with second page

$$E_2^{p,q} = R^p f(R^q g(\mathcal{F}))$$

that abuts to $R^n(fg)(\mathcal{F})$: this is a particular case of the spectral sequence of composition of functors (see [8]). From this, we obtain the formula

$$(fg)_!(\mathcal{F}) = f_!(g_!(\mathcal{F}))$$

whence finally the fact that $(fg)_! = f_!g_!$.

6 Chern classes

The fact that $K_1(X) = K(X)$ allows us to extend the definition of Chern classes to arbitrary coherent sheaves.

First consider the case where the base field is \mathbb{C} ; every vector bundle E over X defines Chern classes $c_i(E) \in H^{2i}(X, \mathbb{Z})$. If we have an exact sequence

$$0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$$

then we know that $c_p(E) = \sum_{r+s=p} c_r(E') \cdot c_s(E'')$.

As in §5.2, this can be understood as a multiplicative property of the Chern polynomial $c_t(E) = \sum c_p(E)t^p$, and allows us to define $c_t(x)$ for all $x \in K(X)$. The $c_p(x)$ are homogeneous elements of degree $2p$ of $H^*(X, \mathbb{Z})$.

In the case of an arbitrary base field, Grothendieck proceeded in the same way, but replacing $H^*(X)$ with the graded ring $A(X)$ of *cycle classes* on X , under linear equivalence (‘{a} la Chow). We recall only that a cycle Z on X of codimension p [i.e. a degree- p element of $A(X)$] is said to be linearly equivalent to zero if there exists a cycle H on $X \times \mathbf{D}$ (where \mathbf{D} denotes the projective line) such that $Z = H_a - H_b$ for points $a, b \in \mathbf{D}$; we denote by H_a the projection to X of $H \cdot (X \times \{a\})$ if it is proper. Chow and Samuel¹ have shown that this equivalence relation possesses all sorts of reasonable properties, and Chow has shown (unpublished) that we could also construct a theory of Chern classes for vector bundles², with these classes being elements of $A(X)$. Note that, even in the classical case, this definition is somehow *finer* than the cohomological definition (since two cycles can indeed be homologous without being linearly equivalent).

| p. 112

In what follows, we denote by $A(X)$ either this ring of cycle classes or $H^*(X)$, and leave it to the reader to choose between the two theories.

Note that, in all cases, if $f: Y \rightarrow X$ is a morphism (resp. a proper morphism), then we can associate to it a homomorphism $f^*: A(X) \rightarrow A(Y)$ (resp. a homomorphism $f_*: A(Y) \rightarrow A(X)$). The formula

$$f_*(y \cdot f^*(x)) = f_*(y) \cdot x$$

holds true.

All of the usual formal constructions explained in the work by Hirzebruch [10] can be applied to the Chern classes $c_p(x)$ of an element $x \in K(X)$. We can, for example, define the *Todd class* $T(x) \in A(X) \otimes \mathbb{Q}$ of an element x : we write $c_t(x)$ formally in the form $\prod(1 + a_i t)$, and let $T(x) = \prod a_i / (1 - e^{-a_i})$. Then

$$T(x + y) = T(x) \cdot T(y).$$

Similarly, we can define the ‘‘exponential’’ Chern class, denoted by $\text{ch}(x)$ (which is also an element of $A(X) \otimes \mathbb{Q}$), by

$$\text{ch}(x) = \text{rank}(x) + \sum (e^{a_i} - 1)$$

where $\text{rank}(x)$ is the *rank* of x [it is the unique homomorphism from $K(X)$ to \mathbb{Z} that sends a vector bundle to its dimension]. Then

$$\text{ch}(x + y) = \text{ch}(x) + \text{ch}(y)$$

$$\text{ch}(x \cdot y) = \text{ch}(x) \cdot \text{ch}(y)$$

¹See Chow [6] and Samuel [11]. See also the *Séminaire Chevalley* [5].

²See the paper by Grothendieck which follows this present work [9].

by the analogous properties of vector bundles. With this, we can calculate $\text{ch}(x)$ in terms of $c_i(x)$ and $\text{rank}(x)$ by “universal” formulas. If $f: Y \rightarrow X$ is a morphism, then

$$c_p(f^!(x)) = f^*(c_p(x)) \text{ch}(f^!(x)) = f^*(\text{ch}(x))$$

for $x \in K(X)$.

These formulas are well known when $x = \gamma(E)$, with E a vector bundle on X , and the general case follows from linearity, by applying [Theorem 2](#).

7 Statement of the Riemann–Roch theorem; First simplifications

Let $f: Y \rightarrow X$ be a proper morphism, with X and Y irreducible non-singular quasi-projective varieties. We denote by $T(X)$ the Todd class of the tangent bundle of X ; it is an element of $A(X) \otimes \mathbb{Q}$. Now let $y \in K(Y)$. Then:

| p. 113

Theorem (Riemann–Roch). —

$$f_*(\text{ch}(y) \cdot T(Y)) = \text{ch}(f_!(y)) \cdot T(X).$$

[The two sides of the equality are thought of as elements of $A(X) \otimes \mathbb{Q}$; with this in mind, we can say that [R–R](#) is a formula “modulo torsion”; Grothendieck has more precise formulas, without torsion — i.e. that hold in $A(X)$ — but he does not yet know how to prove them except in characteristic zero.]

We now show how the [Riemann–Roch theorem](#), in Grothendieck’s form, *implies the Riemann–Roch formula of Hirzebruch* [\[10\]](#):

We apply [R–R](#) to a projective Y , with X consisting of a single point, and y the class of a coherent sheaf \mathcal{F} on Y . Since $A(X)$ is simply \mathbb{Z} in dimension 0, and is zero in higher dimensions, $f_*(u)$, for $u \in A(Y)$, is simply the terms $x_n(u)$ of degree $n = \dim Y$ in u . Also, $T(X) = 1$, and $f_!(y)$ is the alternating sum of the sheaves $R^q f_*(\mathcal{F})$; a sheaf on a point is simply a vector space of finite dimension; in particular, $R^q f_*(\mathcal{F})$ is the vector space $H^q(Y, \mathcal{F})$. On X , the map $\mathcal{G} \mapsto \text{ch}(\mathcal{G})$ consists simply of taking the *rank* of a sheaf; the right-hand side of [R–R](#) thus becomes $\sum (-1)^p \dim H^p(Y, \mathcal{F}) = \chi(Y, \mathcal{F})$, and [R–R](#) then reduces to the form given by Hirzebruch:

$$x_n(\text{ch}(\mathcal{F}) \cdot T(Y)) = \chi(Y, \mathcal{F}).$$

We note that this formula is proven for any coherent sheaf, and not simply for vector bundles; this generality is somewhat illusory, by the linear character of [R–R](#) and by [Theorem 2](#).

The proof of [R–R](#) proceeds by reduction to particular cases of a projection and an injection. For this, we use the following lemma:

Lemma 15.

Let $Z \xrightarrow{g} Y \xrightarrow{f} X$ be proper morphisms. Let $z \in K(Z)$ and let $y = g^!(z)$. Then:

- a. If [R–R](#) is true for $\{g, z\}$ and $\{f, y\}$, then it is also true for $\{fg, z\}$.
- b. If [R–R](#) is true of $\{fg, z\}$ and $\{f, y\}$, and if f_* is injective, then [R–R](#) is also true for $\{g, z\}$.

Proof. First we prove (a). By **R–R** for g , we have

$$g_*(\text{ch}(z) \cdot T(Z)) = \text{ch}(y) \cdot T(Y).$$

Applying f_* to both sides, and taking into account the fact that $(fg)_* = f_*g_*$, we see that | p. 114

$$(fg)_*(\text{ch}(z) \cdot T(Z)) = f_*(\text{ch}(y) \cdot T(Y)).$$

Applying **R–R** for $\{f, y\}$, we see that the right-hand side of the above is equal to $\text{ch}((fg)_!(z) \cdot T(X))$, which indeed proves **R–R** for $\{fg, z\}$.

Now we prove (b). Let

$$\begin{aligned} u &= g_*(\text{ch}(z) \cdot T(Z)) \\ v &= \text{ch}(y) \cdot T(Y). \end{aligned}$$

We wish to show that $u = v$. Given the hypothesis on f_* , it suffices to prove that $f_*(u) = f_*(v)$. But **R–R** for $\{fg, z\}$ proves that

$$f_*(u) = \text{ch}(x) \cdot T(X) \quad \text{where } x = (fg)_!(z) = f_!(y).$$

Similarly, **R–R** for $\{f, y\}$ proves that

$$f_*(v) = \text{ch}(x) \cdot T(X).$$

□

Now let Y and Y' be varieties, and consider their product $Y \times Y'$. The projections $Y \times Y' \rightarrow Y$ and $Y \times Y' \rightarrow Y'$ define homomorphisms $K(Y) \rightarrow K(Y \times Y')$ and $K(Y') \rightarrow K(Y \times Y')$, whence a homomorphism $K(Y) \otimes K(Y') \rightarrow K(Y \times Y')$. By an abuse of language, we also denote by $y \otimes y'$ the image in $K(Y \times Y')$ of the tensor product of two elements $y \in K(Y)$ and $y' \in K(Y')$.

Lemma 16. *Let $f: Y \rightarrow X$ and $f': Y' \rightarrow X'$ be proper morphisms, and let $y \in K(Y)$ and $y' \in K(Y')$. If **R–R** is true for $\{f, y\}$ and $\{f', y'\}$, then it is true for $\{f \times f', y \otimes y'\}$.*

(We denote by $f \times f': Y \times Y' \rightarrow X \times X'$ the product of f and f' .)

Proof. The proof consists of a calculation analogous to that for **Lemma 15**; we have to use the following formulas:

- i. $(f \times f')_!(y \otimes y') = f_!(y) \otimes f'_!(y')$;
- ii. $(f \times f')_*(x \otimes x') = f_*(x) \otimes f'_*(x')$, where $x \in A(Y)$ and $x' \in A(Y')$;
- iii. $\text{ch}(y \otimes y') = \text{ch}(y) \otimes \text{ch}(y')$.

Formula (i) can be proven by taking $y = \gamma(\mathcal{F})$ and $y' = \gamma(\mathcal{F}')$, and applying the Künneth formula to calculate $(f \times f')_!(y \otimes y')$; the validity of the Künneth formula for coherent sheaves follows from the calculation of the cohomology by covers and from the Eilenberg–Zilber theorem (see the book by Godement on sheaves).

Formula (ii) is immediate, whether we consider the point of view of classes of cycles (for linear equivalence), or the cohomological point of view (in the classical case). | p. 115

Formula (iii) is a consequence of the multiplicative property of ch . □

[In fact, we will only use the above lemma in the case where one of the varieties X' and Y' consists of a single point.]

[Lemma 15](#), along with [the Corollary of Proposition 4](#), shows that *it suffices to prove R–R in the two following cases:*

- a. $Y = X \times \mathbf{P}$, with \mathbf{P} a projective space, and $f: X \times \mathbf{P} \rightarrow X$ the projection onto the first factor.
- b. an injection $f: Y \rightarrow X$ of a closed subvariety Y of X .

By [Lemma 16](#), (a) follows from:

- a'. The homomorphism $K(X) \otimes K(\mathbf{P}) \rightarrow K(X \times \mathbf{P})$ is surjective;
- a''. R–R is true for the map from \mathbf{P} to a point (or, in other words, the R–R-Hirzebruch formula is true for \mathbf{P}).

The next two sections are dedicated to the proof of (a') and (a''). The case of an injection (which is more difficult) will be covered later on.

8 Exactness and homotopy properties of $K(X)$

Proposition 7. *Let X be an algebraic variety (singular or not), and X' a closed subvariety of X , and set $U = X \setminus X'$. Then we have an exact sequence*

$$K(X') \rightarrow K(X) \rightarrow K(U) \rightarrow 0.$$

[The homomorphism $K(X') \rightarrow K(X)$ is that given by extending a sheaf on X' by zero outside of X' ; in the case where X and X' are non-singular, and denoting the canonical injection $X' \rightarrow X$ by i , this homomorphism is $i_!$. As for $K(X) \rightarrow K(U)$, this is the restriction homomorphism; in the case where X is non-singular, and denoting the canonical injection $U \rightarrow X$ by j , this homomorphism is $j^!$.]

Proof. Let $A = K(X)/\text{Im}K(X')$. We will define a homomorphism from $K(U)$ to A . For this, let \mathcal{F} be a coherent sheaf on U ; by [Proposition 2](#), we can extend \mathcal{F} to a sheaf \mathcal{G} on X . We will show that the image of $\gamma_X(\mathcal{G})$ in A does not depend on \mathcal{G} , but only on \mathcal{F} . So let \mathcal{G} and \mathcal{G}' be extensions of \mathcal{F} . Embed \mathcal{F} diagonally into $\mathcal{F} \times \mathcal{F} = (\mathcal{G} \times \mathcal{G}')|_U$; by [Proposition 1](#), there exists a subsheaf \mathcal{G}'' of $\mathcal{G} \times \mathcal{G}'$ such that its restriction to U is \mathcal{F} , and everything then relies on proving that

$$\gamma_X(\mathcal{G}'') \equiv \gamma_X(\mathcal{G}') \pmod{\text{Im}K(X')}$$

and similarly for \mathcal{G}' . But, since $\mathcal{G}'' \subset \mathcal{G} \times \mathcal{G}'$, we have a homomorphism $f: \mathcal{G}'' \rightarrow \mathcal{G}$ that is bijective over U . Let \mathcal{N} and \mathcal{Q} be the kernel and cokernel (respectively) of f ; then

$$\gamma_X(\mathcal{G}'') - \gamma_X(\mathcal{G}) \equiv \gamma_X(\mathcal{N}) - \gamma_X(\mathcal{Q}).$$

But \mathcal{N} and \mathcal{Q} are supported outside of X' ; if \mathcal{I} denotes the sheaf of ideals of X' in \mathcal{O}_X , then there exists an integer $n > 0$ such that $\mathcal{I}^n \mathcal{N} = 0$, and similarly for \mathcal{Q} (this is a local statement that is essentially equivalent to the “Nullstellensatz,” see [\[14\]](#)); we

| p. 116

thus conclude that \mathcal{N} and \mathcal{Q} admit composition series whose successive quotients are annihilated by \mathcal{I} , i.e. are coherent on X' , and so

$$\gamma_X(\mathcal{Q}) \equiv \gamma_X(\mathcal{N}) \equiv 0 \pmod{\text{Im}K(X')}.$$

The independence of $\gamma_X(\mathcal{G}) \pmod{\text{Im}K(X)}$ is thus proven, and so we obtain a well-defined element $\eta(\mathcal{F}) \in A$. If $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ is an exact sequence on U , then we can extend \mathcal{F} and \mathcal{G} to X , and \mathcal{F}' to a subsheaf \mathcal{G}' of \mathcal{G} (see [Proposition 1](#)). The sheaf $\mathcal{G}'' = \mathcal{G}/\mathcal{G}'$ extends \mathcal{F}'' , which proves that $\eta(\mathcal{F}) = \eta(\mathcal{F}') + \eta(\mathcal{F}'')$. The operation η , being additive, defines $\eta: K(U) \rightarrow A$. If we denote by ε the canonical homomorphism from A to $K(U)$, we immediately see that $\eta \circ \varepsilon = 1$ and $\varepsilon \circ \eta = 1$, which finishes the proof. \square

Now let X and Y be varieties. Denote by $p: X \times Y \rightarrow X$ the canonical projection. If X and Y are *non-singular*, then the homomorphism $p^1: K(X) \rightarrow K(X \times Y)$ is defined (see [§5.3](#), as well as [§7](#)). In fact, this homomorphism can be defined *in the general case*. This comes from the fact that $\mathcal{O}_{X \times Y}$ is \mathcal{O}_X -flat (i.e. an annihilator of Tor), since it is a ring of fractions of the usual tensor product $\mathcal{O}_X \otimes_k \mathcal{O}_Y$. We can thus set $p^1(\mathcal{F}) = \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_{X \times Y}$ for any coherent sheaf \mathcal{F} on X , and $p^1(\mathcal{F})$ is additive, and so defines some $K(X) \rightarrow K(X \times Y)$. Thus:

Proposition 8. *If Y is an affine line, then the homomorphism $p^1: K(X) \rightarrow K(X \times Y)$ is bijective.*

Proof. Let a be the origin in Y , and identify X with $X \times \{a\} \subset X \times Y$. We have an exact sequence:

$$0 \rightarrow \mathcal{O}_{X \times Y} \xrightarrow{t} \mathcal{O}_{X \times Y} \rightarrow \mathcal{O}_X \rightarrow 0$$

where t denotes the projection $X \times Y \rightarrow Y$, considered as a function to $X \times Y$. This exact sequence shows that $\text{Tor}_p^{\mathcal{O}_{X \times Y}}(\mathcal{O}_X, \mathcal{F}) = 0$ for $p \geq 2$, if \mathcal{F} is a coherent sheaf on $X \times Y$. We can thus define $\pi_a: K(X \times Y) \rightarrow K(X)$ by setting

$$\pi_a(\mathcal{F}) = \text{Tor}_0(\mathcal{O}_X, \mathcal{F}) - \text{Tor}_1(\mathcal{O}_X, \mathcal{F});$$

we see, furthermore, that $\pi_a \circ p^1 = 1$, which already shows that p^1 is *injective*. From now on, we will consider $K(X)$ as *embedded* into $K(X \times Y)$.

To prove that $K(X) = K(X \times Y)$, we argue by induction on $n = \dim X$, and we use the following diagram (where X' is closed in X , and $U = X \setminus X'$):

$$\begin{array}{ccccccc} K(X') & \longrightarrow & K(X) & \longrightarrow & K(U) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ K(X' \times Y) & \longrightarrow & K(X \times Y) & \longrightarrow & K(U \times Y) & \longrightarrow & 0 \end{array}$$

By [Proposition 7](#), the rows of this diagram are exact. We thus conclude that, if $K(X') = K(X' \times Y)$ (which is the case, by the induction hypothesis, if $\dim X' < n$), every element of $K(X \times Y)$ whose restriction to $U \times Y$ is in $K(U)$ is also an element of $K(X)$. In other words, we can “neglect” the subvarieties of X of dimension $< n$. In particular, we can suppose X to be *affine, non-singular, and irreducible* (since the complement of the set of singular points

| p. 117

is a union of disjoint irreducible varieties). We will make use of the following dévissage lemma:

Lemma 17. *Let Z be an algebraic variety. Then the $\gamma_Z(\mathcal{O}_T)$ generate $K(Z)$, where T runs over the set of irreducible subvarieties of Z .*

Proof. We argue by induction on $\dim Z$; using [Proposition 7](#), we can assume that Z is irreducible. Let \mathcal{F} be a sheaf on Z ; if \mathcal{F} is a torsion sheaf, then it is concentrated on a subvariety, and the induction hypothesis shows that it is contained in the subgroup $K'(Z)$ of $K(Z)$ generated by the $\gamma_Z(\mathcal{O}_T)$. In the case where \mathcal{F} is torsion-free, we embed it into $\mathcal{F} \otimes_{\mathcal{O}_Z} K$, where K is the field of rational functions on Z ; we then have that $\mathcal{F} \otimes K = K^n$, and from this we conclude that \mathcal{F} is isomorphic to $(\mathcal{O}_Z)^n$ modulo a torsion sheaf; whence the result. (For more details, see [\[14\]](#), or [\[7\]](#).) \square

By applying [Lemma 17](#) to the case that interests us, we see that it suffices to show that $\gamma(\mathcal{O}_T) \in K(X)$ for every irreducible subvariety T of $X \times Y$. If $\text{proj}_X(T) \neq X$, then it is evident, by the induction hypothesis; if $T = X \times Y$, then it is even more evident. Thus the only remaining case is when $\dim T = n$, and $\text{proj}_X(T)$ is dense in X . If A denotes the coordinate ring of X , then T corresponds to a minimal prime ideal \mathfrak{p} in the coordinate ring $A[t]$ of $X \times Y$; the fact that the projection from T to X is dense implies that $A \cap \mathfrak{p} = 0$. Let S be the set of invertible elements of A , and let $K = A_S$ be the field of fractions of A . We have that $A[t]_S = K[t]$, and the fact that $\mathfrak{p} \cap S = \emptyset$ implies that \mathfrak{p} is of the form $\mathfrak{p}' \cap A[t]$, where \mathfrak{p}' is a non-zero prime ideal of $K[t]$. There thus exists an irreducible polynomial $P(t)$, with coefficients in A if we want, such that \mathfrak{p} is the set of polynomials of $A[t]$ that are divisible by $P(t)$ in $K[t]$. Then

$$A[t] \supset \mathfrak{p} \supset P(t) \cdot A[t] = \mathfrak{q}.$$

The sheaf \mathcal{O}_T corresponds to the module $A[t]/\mathfrak{p}$; let \mathcal{F} be the sheaf that corresponds to $A[t]/\mathfrak{q}$. Since $\mathfrak{p}_S = \mathfrak{q}_S$, there exists some $a \in S$ such that $a \cdot (\mathfrak{p}/\mathfrak{q}) = 0$, which shows that \mathcal{O}_T is congruent to \mathcal{F} modulo an element of $K(X' \times Y)$, with $\dim X' < \dim X$. Also, the exact sequence

$$0 \rightarrow \mathcal{O}_{X \times Y} \xrightarrow{P(t)} \mathcal{O}_{X \times Y} \rightarrow \mathcal{F} \rightarrow 0$$

shows that $\gamma(\mathcal{F}) = 0$ in $K(X \times Y)$, whence

$$\gamma(\mathcal{O}_T) \in \text{Im} K(X' \times Y) = \text{Im} K(X') \subset K(X)$$

which finishes the proof. \square

Remark.[Cartier] We can avoid the need for the dévissage lemma by applying, to an arbitrary module M over $A[t]$, the procedure applied above to the prime ideal \mathfrak{p} . We form $M \otimes_A K = M_S$, which is a $K[t]$ -module. The structure of modules over principal rings shows that the group of classes of $K[t]$ -modules is infinite cyclic, generated by the module $A[t]$. Since passing to M_S consists of “neglecting” everything that is concentrated on $X' \times Y$, the result follows.

| p. 118

Corollary. *If Y is an affine space k^n , then*

$$K(X) = K(X \times Y).$$

Proof. This is immediate, by induction on $\dim Y$. □

9 Proof of the Riemann–Roch theorem for $f: X \times \mathbf{P} \rightarrow X$

We will first prove claim (a') from the end of §7:

Proposition 9. *For every variety X , and every projective space \mathbf{P} , the homomorphism $K(X) \otimes K(\mathbf{P}) \rightarrow K(X \times \mathbf{P})$ is surjective.*

Proof. We argue by induction on $\dim \mathbf{P}$. For $\dim \mathbf{P} = 0$, the proposition is trivial. So let \mathbf{P}' be a hyperplane in \mathbf{P} , and let $U = \mathbf{P} \setminus \mathbf{P}'$. We then have a diagram of exact sequences

$$\begin{array}{ccccccc} K(X) \otimes K(\mathbf{P}') & \longrightarrow & K(X) \otimes K(\mathbf{P}) & \longrightarrow & K(X) \otimes K(U) & \longrightarrow & 0 \\ \downarrow \varepsilon_1 & & \downarrow \varepsilon_2 & & \downarrow \varepsilon_3 & & \\ K(X \times \mathbf{P}') & \longrightarrow & K(X \times \mathbf{P}) & \longrightarrow & K(X \times U) & \longrightarrow & 0 \end{array}$$

This diagram commutes: this is trivial for the second and third squares, and, for the first square, it suffices to verify locally, on affine opens, for example. Given the induction hypothesis, ε_1 is surjective. But U is an affine space; the [Corollary of Proposition 8](#) then shows that $K(U) = \mathbb{Z}$, and that $K(X \times U) = K(X)$, whence the fact that ε_3 is bijective. The five lemma then shows that ε_2 is surjective. □

Remark. The above proof shows that

$$K(X) \otimes K(Y) \rightarrow K(X \times Y)$$

is surjective whenever Y admits an *algebraic cellular decomposition*, where the cells are *affine spaces*. This is notably the case if Y is a Grassmannian.

We now proceed to the proof of claim (a''):

Proposition 10. *Let \mathcal{F} be a coherent sheaf on a projective space \mathbf{P} of dimension r . Then | p. 119*

$$x_r(\text{ch}(\mathcal{F}) \cdot T(\mathbf{P})) = \chi(\mathbf{P}, \mathcal{F})$$

holds true.

Proof. Let H be a hyperplane, and x its class in $A^1(\mathbf{P})$. We know that the Chern polynomial of \mathbf{P} is $(1 + tx)^{r+1}$; thus

$$T(\mathbf{P}) = x^{r+1}/(1 - e^{-x})^{r+1}.$$

We also know that \mathcal{F} corresponds to a graded module M over $k[X_0, \dots, X_r]$. By the Hilbert syzygy theorem, M admits a finite resolution by free graded modules; it follows

that \mathcal{F} is equal [in $K(X)$] to a linear combination of sheaves $\mathcal{O}(n)$, defined in [12], and it suffices to verify the R–R–Hirzebruch formula for $\mathcal{F} = \mathcal{O}(n)$. This sheaf is exactly the sheaf associated to the divisor nH ; thus $\text{ch}(\mathcal{F}) = e^{nx}$. Now, a direct calculation shows that $\chi(\mathbf{P}, \mathcal{F})$ is equal to $\binom{n+r}{r}$ (see [12]), and we are thus led to prove the formula:

$$x_r[e^{nx} \cdot x^{r+1}/(1 - e^{-x})^{r+1}] = \binom{n+r}{r}. \quad (\star)$$

It is useful to write this formula in terms of residues:

$$\text{Res}[e^{nx} dx/(1 - e^{-x})^{r+1}] = \binom{n+r}{r}. \quad (\star\star)$$

By defining the new variable $y = 1 - e^{-x}$, we see that the left-hand side is equal to

$$\begin{aligned} \text{Res}(dy \cdot y^{-r-1} \cdot (1-y)^{-n-1}) &= z_r((1-y)^{-n-1}) \\ &= (-1)^r \binom{-n-1}{r} = \binom{n+r}{r}. \end{aligned}$$

□

Since we have proven (a') and (a''), we can state:

Corollary. *R–R is true for the projection $X \times \mathbf{P} \rightarrow X$.*

Remark. The fact that $K(\mathbf{P})$ is generated by the $\mathcal{O}(n)$ can also be seen, without using the syzygy theorem, by the cellular decomposition of \mathbf{P} . We again obtain the complete structure of $K(\mathbf{P})$: if we denote by α the class in $K(\mathbf{P})$ of the \mathcal{O}_H , then the elements $1, \alpha, \alpha^2, \dots, \alpha^r$ form a *basis* of $K(\mathbf{P})$, and $\alpha^{r+1} = 0$. The $\mathcal{O}(n)$ are equal to $(1 - \alpha)^{-n}$, as shown by the exact sequence

$$0 \rightarrow \mathcal{O}(-1) \rightarrow \mathcal{O} \rightarrow \mathcal{O}_H \rightarrow 0.$$

10 General remarks on the injection of a subvariety

10.1 Notation

Before continuing the proof of R–R, we first discuss the local resolution of the structure sheaf of a subvariety, and study some of its consequences. Some of these will be more general than necessary for our purposes, but they are interesting in their own right, and are no more difficult than the particular cases that we will need. | p. 120

In what follows, Y is a (non-singular) subvariety of X , i the injection of Y into X , p the codimension of Y in X , E the normal bundle of Y , and $\mathcal{I}(Y)$ the sheaf of local ideals of Y . We thus have the exact sequence

$$0 \rightarrow \mathcal{I}(Y) \rightarrow \mathcal{O}_X \xrightarrow{r} \mathcal{O}_Y \rightarrow 0 \quad (1)$$

where r is the restriction.

Finally, write F^* to mean the dual of a vector bundle F , and $[Z]$ to mean the bundle associated to a divisor Z . Recall that $c([Z]) = 1 + Z$.

10.2 Local parameters; Normal bundle

Let $a \in Y$. Then there exists an affine open subset $U \subset X$ containing a , and functions f_1, \dots, f_p that are regular on U , and that form a system of local parameters, or a system of “uniformising” parameters, for Y . This implies that $Y \cap U$ is defined by the equations $f_1 = \dots = f_p = 0$, that df_1, \dots, df_p are linearly independent at every point in $Y \cap U$, that the f_i form a basis for the ideal $\mathcal{I}(Y)_b$ of Y in $\mathcal{O}_{b,X}$ ($b \in Y \cap U$), and, finally, that this ideal is “perfect,” i.e. that the annihilator of f_i in $\mathcal{O}_{b,X}/(f_1, \dots, f_{i-1})$ is zero ($1 \leq i \leq p$; $f_0 = 0$).

At every point $b \in Y \cap U$, the df_i form a basis for E_b^* . Also, $\mathcal{I}(Y)/\mathcal{I}(Y)^2$ is a sheaf concentrated on Y , and annihilated by $\mathcal{I}(Y)$, and is thus a sheaf of \mathcal{O}_Y -modules. It is immediate that $f \mapsto df$ is a homomorphism of \mathcal{O}_Y -modules from $\mathcal{I}(Y)/\mathcal{I}(Y)^2$ to the sheaf $\mathcal{O}_Y(E^*)$ of germs of sections of E^* . By using the local parameters, we see that

the map $D: \mathcal{I}(Y)/\mathcal{I}(Y)^2 \rightarrow \mathcal{O}_Y(E^)$ defined by $f \mapsto df$ is an isomorphism of \mathcal{O}_Y -modules.*

10.3 Local resolution of \mathcal{O}_Y over \mathcal{O}_X

Let V be a vector space over k of dimension p , and e_1, \dots, e_p be a basis of V , and define $\mathfrak{M}_i = \mathcal{O}_X \otimes_k \wedge^i V$. We endow the direct sum \mathfrak{M} of the \mathfrak{M}_i with a differential d defined by³

$$d(f \otimes e_{i_1} \wedge \dots \wedge e_{i_k}) = \sum_j (-1)^j f \cdot f_{i_j} \otimes e_{i_1} \wedge \dots \wedge \widehat{e_{i_j}} \wedge \dots \wedge e_{i_k}.$$

Then the sequence

$$0 \rightarrow \mathfrak{M}_p \xrightarrow{d} \mathfrak{M}_{p-1} \rightarrow \dots \rightarrow \mathfrak{M}_1 \xrightarrow{d} \mathcal{O}_X \xrightarrow{r} \mathcal{O}_Y \rightarrow 0 \quad (2)$$

is exact at every point of U , and \mathfrak{M} is thus, on U , a projective resolution of \mathcal{O}_Y over \mathcal{O}_X . The exactness of (2) at a point $b \in U \cap Y$ is a well-known result (see, for example, [3, Proposition 4.3, p. 151], taking into account the fact that $\mathcal{O}_{b,Y} = \mathcal{O}_{b,X}/(f_1, \dots, f_p)$). For $b \in U \setminus (U \cap Y)$, we have that $\mathcal{O}_{b,Y} = 0$, and the exactness of (2) is then an elementary exercise left to the reader. | p. 121

Remark. If Y is a divisor, then (1) is exactly another way of writing the exact sequence of sheaves (which is exact on any X)

$$0 \rightarrow \mathcal{O}_X([Y]^{-1}) \xrightarrow{f} \mathcal{O}_X \rightarrow \mathcal{O}_Y \rightarrow 0, \quad (3)$$

where f is the multiplication by a local equation on Y . This exact sequence thus shows that $\gamma(Y) = 1 - [Y]^{-1}$.

Proposition 11. *Let Y_1, \dots, Y_m be non-singular subvarieties of X such that Y_i intersects transversally with*

$$Y_{i-1} \cap \dots \cap Y_1 \quad (\text{for } i = 2, \dots, m).$$

Then

$$\gamma(Y_1 \cap \dots \cap Y_m) = \prod_i \gamma(Y_i).$$

³As usual, the notation $\widehat{}$ indicates that the symbol below the hat should be omitted.

Proof. An evident induction on m shows that it suffices to consider the case of two subvarieties Y and Z that intersect transversally.

By definition, $\gamma(Y) \cdot \gamma(Z)$ is the alternating sum of the $\text{Tor}_i^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_Z)$. It is clear that these sheaves, and the sheaf $\mathcal{O}_{Y \cap Z}$, are equal (and zero) at any point not contained inside $Y \cap Z$, which, in particular, proves our claim if $Y \cap Z = \emptyset$.

Now let $a \in Y \cap Z$. Since Y and Z intersect transversally, we can find an affine open subset U containing a , and functions f_1, \dots, f_p and g_1, \dots, g_q that are regular on U such that the f_i (resp. the g_j ; resp. the f_i and the g_j) form a system of uniformising parameters for Y (resp. for Z ; resp. for $Y \cap Z$). To calculate the $\text{Tor}_i(\mathcal{O}_Y, \mathcal{O}_Z)$, we resolve \mathcal{O}_Y by using (2). We must thus determine the homology of the complex

$$0 \rightarrow \mathfrak{M}_n \otimes_{\mathcal{O}_X} \mathcal{O}_Z \rightarrow \dots \rightarrow \mathcal{O}_X \otimes_{\mathcal{O}_X} \mathcal{O}_Z$$

which can also be written as

$$0 \rightarrow \mathcal{O}_Z \otimes_k \bigwedge^p V \rightarrow \dots \rightarrow \mathcal{O}_Z \otimes_k V \rightarrow \mathcal{O}_Z$$

endowed with the same differential as above, with the f_i being considered as element of \mathcal{O}_Z . Since $\mathcal{O}_{a,Z} = \mathcal{O}_{a,X}/(g_1, \dots, g_q)$, the f_i generate a perfect ideal inside; so this complex is acyclic (see §10.3) and its 0-th homology group is $\mathcal{O}_{a,Z}(f_1, \dots, f_p)$, i.e. $\mathcal{O}_{a,Y \cap Z}$. Thus

$$\text{Tor}_0 = \mathcal{O}_{Y \cap Z} \quad \text{and} \quad \text{Tor}_i = 0 \quad (\text{for } i \geq 1),$$

which finishes the proof. □

| p. 122

Corollary. *Let Y be a non-singular hyperplane section of Z , and let k be the dimension of X . Then $(1 - [Y])^{k+1} = 0$.*

Proof. We take the non-singular hyperplane sections Y_1, \dots, Y_{k+1} whose intersection is empty, and such that the Y_i intersect transversally with $Y_{i-1} \cap \dots \cap Y_1$ ($i = 2, \dots, k+1$). Since $[Y_i] = [Y]$, Proposition 11 and the preceding remark show that $(1 - [Y]^{-1})^{k+1} = 0$, whence the corollary, since $[Y]$ is invertible in $K(X)$. □

10.4 —

We will use, both here and later on, the following remark; let \mathcal{F} and \mathcal{G} be sheaves of algebras on X . Then the direct sum of the $\text{Tor}_i^{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})$ canonically admits the structure of a sheaf of graded (by i) algebras. If \mathcal{F} admits, on U , a resolution \mathcal{L} over \mathcal{O}_X that is a sheaf of graded algebras, then the product in the homology of $\mathcal{L} \otimes \mathcal{G}$ induced by the products in \mathcal{L} and \mathcal{G} coincides, on U , with the the aforementioned coupling of the Tor_i . [Details left to the read; see [3, Chap. IX] for some analogous and more general considerations. This follows from the classical result on maps from an acyclic complex ([3, Proposition 11, p. 76]).]

Proposition 12. *We have that $i^! i_!(y) = y \cdot \lambda_{-1}(E^*)$ for all $y \in K(Y)$. In particular, $i^! \gamma(Y) = \lambda_{-1}(E^*)$.*

Proof. By linearity, and [Theorem 2](#), we can restrict to proving [Proposition 12](#) in the case where $y = \mathcal{F}$ is a locally free sheaf. By definition, $i^!_{i_1}(\mathcal{F})$ is the alternating sum of the $\mathrm{Tor}_i^{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_Y)$, and, since \mathcal{F} and the $\lambda^i E^*$ are locally free, $\mathcal{F} \cdot \lambda_{-1} E^*$ is the alternating sum of the $\mathcal{F} \otimes_{\mathcal{O}_Y} \lambda^i E^*$. Recalling [§10.2](#), it thus suffices, to prove [Proposition 12](#), to show that

$$\mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_Y) = \mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{I}(Y) / \mathcal{I}(Y)^2 \quad (4)$$

$$\mathrm{Tor}_i^{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_Y) = \mathcal{F} \otimes_{\mathcal{O}_Y} \lambda^i(\mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_Y)). \quad (5)$$

[In (5), λ^i is the i th exterior power of a sheaf which is locally free, given (4).]

Unless otherwise mentioned, Tor and \otimes are taken over \mathcal{O}_X . The exact sequence of Tor , applied to (1), gives the exact sequence

$$0 \rightarrow \mathrm{Tor}_1(\mathcal{F}, \mathcal{O}_Y) \rightarrow \mathcal{I}(Y) \otimes \mathcal{F} \xrightarrow{g} \mathcal{F}$$

where g is defined by $g(u \otimes v) = u \cdot v$. Since $\mathcal{I}(Y)$ annihilates \mathcal{F} , g has a zero image, and so

$$\mathrm{Tor}_1(\mathcal{F}, \mathcal{O}_Y) = \mathcal{I}(Y) \otimes \mathcal{F}.$$

But, since \mathcal{F} is a sheaf on Y , it is annihilated by $\mathcal{I}(Y)$, and so the image of $\mathcal{I}(Y)^2 \otimes \mathcal{F}$ in $\mathcal{I}(Y) \otimes \mathcal{F}$ is zero, and $\mathcal{I}(Y) \otimes \mathcal{F}$ can be identified with $\mathcal{I}(Y) / \mathcal{I}(Y)^2 \otimes \mathcal{F}$. Finally, since, in this product, the two sheaves are over \mathcal{O}_Y , their \otimes over \mathcal{O} can be identified with their tensor product over \mathcal{O}_Y , which finishes the proof of (4).

Using the notation of [§10.2](#), the $\mathrm{Tor}_i(\mathcal{F}, \mathcal{O}_Y)$ are, on U , the homology groups of the complex

$$0 \rightarrow \mathfrak{M}_p \otimes \mathcal{F} \rightarrow \mathfrak{M}_{p-1} \otimes \mathcal{F} \rightarrow \dots \rightarrow \mathcal{F}$$

which can also be written as

$$0 \rightarrow \mathcal{F} \otimes_k \bigwedge^p V \rightarrow \mathcal{F} \otimes_k \bigwedge^{p-1} V \rightarrow \dots \rightarrow \mathcal{F} \quad (6)$$

endowed with the differential that extends d and is zero on \mathcal{F} . Since the f_i are local sections of $\mathcal{I}(Y)$, and this sheaf annihilates \mathcal{F} , it follows that, in (6), the differential is identically zero, and so

$$\mathrm{Tor}_i(\mathcal{F}, \mathcal{O}_Y) = \mathcal{F} \otimes_k \bigwedge^i V = \mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{O}_Y \otimes_k \bigwedge^i V \quad (\text{for } i \geq 0).$$

In the particular case where $\mathcal{F} = \mathcal{O}_Y$, this formula shows that

$$\mathrm{Tor}_1(\mathcal{O}_Y, \mathcal{O}_Y) = \mathcal{O}_Y \otimes_k V$$

whence we have an isomorphism (for the moment, local, defined by the use of the parameters f_i) of the two sides of (5). But \mathfrak{M} , thought of as a tensor product of the algebras \mathcal{O}_Y and $\bigwedge V$, is clearly a differential graded algebra, and so the remark made at the start of [§10.4](#) applies, and shows that, in fact, this isomorphism is intrinsic (and, in particular, does not depend on the chosen system of local parameters) and holds true over X . \square

10.5 Particular case of the divisor

In this case, we can use the resolution (3), which is global. The $\text{Tor}_i(\mathcal{F}, \mathcal{O}_Y)$ are thus the homology groups of the complex

$$\dots \rightarrow 0 \rightarrow \mathcal{O}_Y([Y]^{-1}) \otimes \mathcal{F} \xrightarrow{f} \mathcal{F}$$

where \mathcal{F} is the multiplication by a local equation; since \mathcal{F} is annihilated by $\mathcal{I}(Y)$, the homomorphism f has a zero image, which implies that

$$\begin{aligned} \text{Tor}_0(\mathcal{F}, \mathcal{O}_Y) &= \mathcal{F} \\ \text{Tor}_1(\mathcal{F}, \mathcal{O}_Y) &= [Y]^{-1} \cdot \mathcal{F} \\ \text{Tor}_i(\mathcal{F}, \mathcal{O}_Y) &= 0 \quad (\text{for } i \geq 2). \end{aligned}$$

By comparing with [Proposition 12](#), we see that the restrictions of $[Y]^{-1}$ to Y and to E^* define the same element of $K(Y)$. In fact,

Proposition 13.

Suppose that $\text{codim } Y = 1$, and denote by L the restriction of $[Y]$ to Y . Then:

- $L = E$;
- $\gamma(Y) = 1 - [Y]^{-1}$;
- $i^! i_!(y) = y \cdot (1 - L^*)$.

| p. 124

Proof. Claim (c) was proven above, and (b) was proven in the remark of [§10.2](#). It remains only to prove (a).

Let (U_i) be a cover of U such that, on U_i , Y is defined by one equation $f_i = 0$, with the differential df_i being non-zero at every point of $Y \cap U_i$, and let $f_{ij} = f_i/f_j$ on $U_i \cap U_j$. We know that $[Y]$ is defined by the system $\{f_{ij}\}$ of transition functions. Also, on U_i , we can identify E^* with the product $U_i \times k$ by applying df_i to the unit section. But, on $Y \cap U_i \cap U_j$, we have that $df_i = f_{ij} \cdot df_j$ (since $f_j = 0$ on Y), and so we can define E^* by the system of transition functions $\{f_{ij}^{-1}\}$, which proves (a). \square

11 Proof of R–R in the particular case of an injection

Given the results of [§7](#) and [§9](#), it remains only to prove R–R for an injection. In the notation of [§10.1](#), the formula to prove is then equivalent to

$$\text{ch } i_! y = i_*(\text{ch } y \cdot T(E)^{-1}) \quad (\text{for } y \in K(Y)). \quad (1)$$

Indeed, E is the quotient of the tangent bundle of X (restricted to Y) by the tangent bundle of Y , so $i^* T(X) = T(Y) \cdot T(E)$, whence

$$\begin{aligned} i_*(\text{ch } y T(E)^{-1}) &= i_*(\text{ch } y \cdot T(Y) \cdot i^*(T(X))^{-1}) \\ i_*(\text{ch } y T(E)^{-1}) &= i_*(\text{ch } y \cdot T(Y)) \cdot T(X)^{-1} \end{aligned}$$

and so (1) gives R–R if we multiply both sides by $T(X)$.

To prove (1), Grothendieck first considers the case where Y is a divisor and $y \in i^!(K(X))$, and then reduces to this case by blowing up X along Y . This first particular case will be covered below, and the general case of the injection will be covered in §13, where we will assume certain properties of blow ups that will be proven in §14, §15, and §16.

We will use, without explanation, the formulas

$$\begin{aligned} f_!(y \cdot f^!(x)) &= f_!(y) \cdot x \\ f_*(y \cdot f^*(x)) &= f_*(y) \cdot x \end{aligned} \tag{2}$$

where the first was proven in §5.3, and the second was mentioned in §6 (whose notation we also continue to use).

Proposition 14. *The equality (1) is true if $\text{codim}_X Y = 1$ and $y = i^!(x)$ ($x \in K(X)$).*

Proof. By using (2) and Proposition 13, we see that

$$\text{ch}(i_! i^!(x)) = \text{ch}(x \cdot i_!(1)) = \text{ch}(x \cdot (1 - [Y]^{-1})).$$

Since $x \mapsto \text{ch } x$ is a ring homomorphism, and since $c([Y]) = 1 + Y$, this implies that

| p. 125

$$\text{ch}(i_!) = \text{ch } x \cdot \text{ch}(1 - [Y]^{-1}) = \text{ch } x \cdot (1 - e^{-Y}).$$

The right-hand side of (1) is

$$\begin{aligned} i_*(\text{ch}(i^! x) \cdot T(L)^{-1}) &= i_*(i^*(\text{ch } x) \cdot i^*(T([Y])^{-1})) \\ &= \text{ch } x \cdot T([Y])^{-1} \cdot i_*(1) \\ &= \text{ch } x \cdot T([Y])^{-1} \cdot Y \end{aligned}$$

and (1) then follows from $T([Y]) = Y/(1 - e^{-Y})$. \square

Corollary 1. *R–R is true if $X = Y \times \mathbf{P}$, where \mathbf{P} is a projective space, and i is the map $a \mapsto (a, p_0)$, with p_0 some fixed point of \mathbf{P} .*

Proof. The map i is the product of the identity on Y with the injection of a point into \mathbf{P} . Given Lemma 16, it suffices to prove R–R in this latter case. If Y is a point, then $K(Y) = \mathbb{Z}$, and it suffices to consider the case where $y = 1$. Since $1 \in i^!(\mathbf{P})$, our claim follows from Proposition 14 if $\dim \mathbf{P} = 1$. Arguing by induction, we can assume that our claim is true for the injection $u: Y \rightarrow H$ of Y into a hyperplane of \mathbf{P} ; given part (a) of Lemma 15, it suffices to prove that R–R is true for $u_!(1)$ and the injection $v: H \rightarrow \mathbf{P}$; this, itself, follows from Proposition 14 if we know that $u_!(1) \in v^!(K(\mathbf{P}))$. But let Z be a second hyperplane and D a line of H such that $Y = D \cap Z \cap H$. Then (Proposition 11) $\gamma_H(Y) = \gamma_H(D) \cdot \gamma_H(H \cap Z)$. By part (b) of Proposition 13, $\gamma_H(H \cap Z) = 1 - [H \cap Z]^{-1}$; since $[H \cap Z]$ can be identified with the restriction of $[H]$ to H , part (c) of Proposition 13 then shows that

$$u_!(1) = \gamma_H(Y) = v^! v_!(\gamma_H(D)) = v^!(\gamma_{\mathbf{P}}(D)).$$

\square

Corollary 2. *If the equality (1) is true whenever $2\dim Y \leq \dim X - 2$, then it is true in general.*

Proof. It suffices to compose i with an injection $X \rightarrow X \times \mathbf{P}$, where \mathbf{P} is of large dimension, and then apply Corollary 1 and part (b) of Lemma 15. \square

12 Blowing up along a subvariety

12.1 Notation

We will write X' to mean the variety obtained by blowing up X along Y , f to mean the projection from X' to X , g to mean the restriction of f to $Y' = f^{-1}(Y)$, and j to mean the injection from Y' to X' . We will thus have the commutative diagram

$$\begin{array}{ccc} Y' & \xrightarrow{j} & X' \\ g \downarrow & & \downarrow f \\ Y & \xrightarrow{i} & X \end{array} \quad (1)$$

As before, E is the normal bundle of Y in X , and $p = \text{codim}_X Y$. We will sometimes write E' to mean the inverse image of E . We will see that f is an isomorphism outside of Y' , that g is the projection into a fibration with fibre equal to the projective space \mathbf{P}_{p-1} of dimension $(p-1)$, that $\text{codim}_{X'} Y' = 1$, and that E' contains the restriction of $[Y']$ to Y' , which will be denoted L . Finally, we let $F = E'/L$, which is a vector bundle of rank $(p-1)$. | p. 126

12.2 Definition of X' by local charts

Over an affine open subset U that does not meet Y , we take U itself as a local chart. Now suppose that $U \cap Y \neq \emptyset$, and that Y admits, in U , uniformising parameters f_i ($1 \leq i \leq p$). Let t_i be homogeneous coordinates of \mathbf{P}_{p-1} . Then $f^{-1}(U)$ is the subvariety of $\mathbf{P}_{p-1} \times U$ defined by

$$U' = f^{-1}(U) = \{(t, u) \mid t_i f_j(u) - t_j f_i(u) = 0\}$$

(with the t_i being the homogeneous coordinates of t) and f is the projection onto the second factor. It is evident that f is a morphism, that it is an isomorphism outside of $f^{-1}(Y \cap U)$, and that, if $u \in Y$, then $f^{-1}(u) = \mathbf{P}_{p-1}$. The variety U' is the union of affine open subsets U'_i , where U'_i is the set of points (t, u) such that $t_i \neq 0$. If we let $f'_j = f_j \circ f$, then the local equation of Y' in U'_i is $f'_i = 0$, and df'_i is a basis for the bundle L at each point of U'_i (see the proof of Proposition 13).

The differentials df_i are linearly independent at every point of $Y \cap U$, and, on $Y \cap U$, we will identify E^* with $U \times k^p$, by applying df_1, \dots, df_p to the canonical basis of k^p . Let $b = (t_1, \dots, t_p) \in g^{-1}(a)$ for $a \in Y \cap U$, and suppose that $t_i \neq 0$. Then the equations in $\mathbf{P} \times U$ of the plane tangent to Y at b are $df_j = (t_j/t_i) df_i$; it follows that f sends, bijectively, the normal space of Y at b to the line $t_1 X_1 + \dots + t_p X_p$, where (X_i) is the dual basis of (df_j) , whence we have a bijection μ_U from $g^{-1}(a)$ to the projective space associated to E_a , and

an inclusion of L into E' , given by sending the fibre L_b of L to the line of E'_b that maps to $t_1X_1 + \dots + t_pX_p$ under the canonical identification of E'_b with E_a .

Let V be a second affine open subset of X that meets Y , and in which Y has local parameters g_1, \dots, g_p . Let \mathbf{P}' be a projective space with homogeneous coordinates s_1, \dots, s_p . Then

$$V' = f^{-1}(V) = \{(s, v) \mid s_i g_j(v) - s_j g_i(v) = 0\}$$

and we will have, as before, a canonical map μ_V from $g^{-1}(a)$ to the projective space of E_a . This leads us to define the change of charts in $U' \times V'$ by the rule:

$$(t, u) = (s, v) \quad \text{if } [v = u \notin Y] \text{ or } [v = u \in Y \text{ and } \mu_U(t) = \mu_V(s)].$$

We must show that this correspondence is an isomorphism. Let $c \in U \cap V$. Since the (f_i) and the (g_i) are two systems of local parameters for Y , there exist elements $a_{ij} \in \mathcal{O}_{c,X}$ that form an invertible matrix in $\mathcal{O}_{c,X}$ such that

$$f_i = \sum_j a_{ij} g_j$$

which also implies that $df_i = \sum_j a_{ij} dg_j$ on Y . In the neighbourhood of a point $c' \in f^{-1}(c)$, the above correspondence is then defined by $(s, v) \mapsto (t, u)$, with $v = u$ and $t_i = \sum a_{ij}(u) s_j$. We see that

$$f_i t_j - f_j t_i = \sum_{m,n} a_{im} a_{jn} (g_m s_n - g_n s_m)$$

and it easily follows that the correspondence in question is an isomorphism.

12.3 Projective embedding of X'

We consider an embedding of X into some projective space \mathbf{P}_N of dimension N , with homogeneous coordinates (z_i) . Let φ_i ($1 \leq i \leq s$) be a basis for the ideal of Y consisting of homogeneous polynomials, and let m be an integer \geq the maximum of the degrees of the φ_i . Let (h_1, \dots, h_M) be a basis for the k -vector space of forms of degree m in the z_i that are zero on Y . We can then take h_i to be the products of the form $\varphi_j \cdot \mu$ ($1 \leq j \leq s$) where μ runs over the set of monomials in the z_i whose degrees are equal to $m - \deg(\varphi_j)$. It follows immediately that:

- i. Given $x \in X \setminus Y$, there exists some j such that $h_j(x) \neq 0$. Given $y \in Y$, there exists some form g of degree m , and a set I of p elements of $\{0, 1, \dots, M\}$ such that the h_i/g ($i \in I$) form, in a suitable affine open neighbourhood of y , a system of local parameters for Y , and such that the $h_i/g \in \mathcal{O}_{y,X}$ ($i = 0, \dots, M$).

The h_i are sections of the bundle H^m , where H is the bundle associated to a hyperplane section of X . We denote by h'_i the corresponding sections of the inverse-image bundle, and we consider, in the usual way, "the map" $h: x' \mapsto (h'_0(x'), \dots, h'_M(x'))$ from X to \mathbf{P}_M defined by this space of sections. *A priori*, this is not really a map, since it is not defined at the points where all the h'_i vanish simultaneously. However, we want to show that:

- ii. h is a morphism;
- iii. if $u', v' \in Y'$ have the same image in Y , and are distinct, then $h(u') \neq h(v')$.

To prove (ii), it suffices to show that, given $x' \in X'$, there exists some index k such that $h'_i/h'_k \in \mathcal{O}_{x',X}$ ($0 \leq i \leq M$), since the $h'_i(x')/h'_k(x')$ will then be the coordinates of $h(x')$, and at least one of these quotients is non-zero. Given (i), the existence of h'_i is evident if $x' \notin Y$. Now let $x' \in Y$. We can apply the second claim of (i), and, to simplify notation, we suppose that the $f_i = h_i/g$ ($i = 1, \dots, p$) form a system of local parameters; let t_1, \dots, t_p be the homogeneous coordinates of x' in \mathbf{P} (the notation from the start of §12.2, and suppose that $t_k \neq 0$). Then $f'_k = 0$ is a local equation for Y' , and so the h'_j/g' are all divisible by f'_k in $\mathcal{O}_{x',X'}$, and so h is a morphism in x' . We also see that the i th homogeneous coordinate of $h(x')$ is t_i/t_k ($1 \leq i \leq p$), which evidently implies (iii).

| p. 128

With this in mind, consider the map Ψ from X' to $X \times \mathbf{P}_M$ defined by $\Psi(x') = (f(x'), (x'))$. Given (ii) and (iii), Ψ is a bijective morphism, and it is clear that Ψ is an isomorphism outside of Y' . By the projective normalisation and the Main Theorem of Zariski, we thus deduce that X' is quasi-projective.

13 Finishing the proof of R–R

Lemma 18. *Let G be a vector bundle of rank k on a variety X . Then*

$$\text{ch}(\lambda_{-1}G) = c_k(G^*)T(G^*)^{-1}.$$

Proof. Write the Chern class $c(G)$ of G in the form

$$c(G) = \prod_{i=1}^k (1 + a_i).$$

Then (see [10])

$$\begin{aligned} c(G^*) &= \prod (1 - a_i), \\ c(\wedge^s G) &= \prod_{i_1 < \dots < i_s} (1 + a_{i_1} + \dots + a_{i_s}), \end{aligned}$$

whence

$$\begin{aligned} c_k(G^*) &= (-1)^k a_1 \dots a_k, \\ \text{ch}(\lambda_{-1}G) &= \prod (1 - e^{a_i}), \end{aligned}$$

and so

$$\text{ch}(\lambda_{-1}G) = c_k(G^*)T(G^*)^{-1}.$$

□

Lemma 19.

In the notation of §12.1, we have:

- a. $f_*(1) = 1$, and so f_*f^* is the identity;
- b. $g_*(c_{p-1}(F)) = 1$;
- c. $f^!i_!(y) = j_!(g^!(y) \cdot \lambda_{-1}F^*) [y \in K(Y)]$;

d. $\lambda_{-1}F \equiv 0 \pmod{(1-L^*)}$ if $p \geq \dim Y + 2$.

Proof. This lemma will be proven in §14, §15, and §16. □

| p. 129

Proposition 15. *R–R is true for an injection.*

Proof. We must (see §11) establish the equality

$$\text{ch } i_!(y) = i_*(\text{ch } y \cdot T(E)^{-1}), \quad (1)$$

and it suffices to do this in the case where $p \geq \dim Y + 2$ (Corollary 2 of Proposition 14). In this case, $g^!(y) \cdot \lambda_{-1}F^* \equiv 0 \pmod{(1-L^*)}$, by part (d) of Lemma 19, and so is in the image of $K(X')$ under $j^!$ (Proposition 13), and (Proposition 14) we can apply R–R to $g^!(y) \cdot \lambda_{-1}F^*$ and j . This gives

$$\text{ch } j_!(g^!y \cdot \lambda_{-1}F^*) = j_*(\text{ch}(g^!y \cdot \lambda_{-1}F^*) \cdot T(L)^{-1}). \quad (2)$$

To deduce (1) from this, it evidently suffices to show that

$$f_*(\text{ch } j_!(g^!y \cdot \lambda_{-1}F^*)) = \text{ch } i_!y \quad (3)$$

$$f_*j_*(\text{ch}(g^!y \cdot \lambda_{-1}F^*) \cdot T(L)^{-1}) = i_*(\text{ch } y \cdot T(E)^{-1}). \quad (4)$$

Part (c) of Lemma 19 shows that the left-hand side of (3) is equal to

$$f_*(\text{ch } f^!i_!y) = f_*^*(\text{ch } i_!y)$$

and thus equal to the right-hand side of (3), given part (a) of Lemma 19.

Since ch is multiplicative, we have that

$$\text{ch}(g^!y \cdot \lambda_{-1}F^*) = \text{ch } g^!y \cdot \text{ch } \lambda_{-1}F^* = g^*(\text{ch } y) \cdot \text{ch } \lambda_{-1}F^*$$

and so (Lemma 18)

$$\text{ch}(g^!y \cdot \lambda_{-1}F^*) = g^*(\text{ch } y)c_{p-1}(F) \cdot T(F)^{-1}.$$

But $E'/L = F$, so

$$g^*(T(E)) = T(E') = T(F) \cdot T(L)$$

whence

$$\text{ch}(g^!y \cdot \lambda_{-1}F^*) \cdot T(L)^{-1} = c_{p-1}(F) \cdot g^*(\text{ch } y \cdot T(E)^{-1}).$$

Given (2) (in §11) and part (b) of Lemma 19, the image under g_* of the right-hand side is equal to $\text{ch } y \cdot T(E)^{-1}$, and the equality in (4) then follows from the fact that $f_*j_* = i_*g_*$. □

14 Proof of parts (a) and (b) of Lemma 19

The map f is an isomorphism outside of Y' , and so is of local degree 1, and so sends a fundamental cycle to a fundamental cycle, whence part (a) of Lemma 19.

The map g_* lowers the degree (which is the geometric codimension) by $p-1$, and corresponds to integration on the fibre. But also, the restriction of L to a fibre \mathbf{P}_{p-1} of g can be thought of as $k^p \setminus \{0\}$, the principal k^* -bundle, on the base \mathbf{P}_{p-1} . We know that the first Chern class of this latter bundle is the opposite of the class of a hyperplane. If we set $-u = c_1(L)$, then this implies that

$$g_*(u^{p-1}) = 1,$$

and we evidently have that $g_*(u^i) = 0$ ($0 \leq i < p-1$) for dimension reasons.

Since $E'/L = F$, we have that

$$\begin{aligned} c(E') &= c(F)(1-u) \\ c(F) &= g^*(c(E)) \cdot (1+u+u^2+\dots) \\ c_{p-1}(F) &= u^{p-1} + g^*(c_1(E)) \cdot u^{p-2} + \dots + g^*(c_{p-1}(E)) \\ g_*(c_{p-1}(F)) &= g_*(u^{p-1}) + c_1(E)g_*(u^{p-2}) + \dots + c_{p-1}(E)g_*(1) \end{aligned}$$

whence

$$g_*(c_{p-1}(F)) = 1.$$

15 Proof of part (c) of Lemma 19

In this section, we will write \mathcal{I} to mean $\mathcal{I}(Y)$, and \mathcal{I}' to mean $\mathcal{I}(Y')$. As we remarked in §10.2, $\mathcal{I}'/\mathcal{I}'^2$ (resp. $\mathcal{I}/\mathcal{I}^2$) is the sheaf of germs of sections of L^* (resp. E^*), and so $\mathcal{I}'/\mathcal{I}'^2 \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'}$ is the sheaf of germs of sections of $g^*(E^*) = E'^*$.

By associating, to an element $u \in \mathcal{I}_x$, the element $u \circ f \in \mathcal{I}'_{x'}(f(x') = x)$, we evidently define a surjective homomorphism of $\mathcal{O}_{Y'}$ -modules $\mu: \mathcal{I}'/\mathcal{I}'^2 \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} \rightarrow \mathcal{I}'/\mathcal{I}'^2$. This again shows that E'^* is sent to L^* , and thus that L injects into E' ; the kernel of μ is the sheaf $\mathcal{O}_{Y'}(F^*)$ of germs of sections of F^* , as we can easily verify; in fact, this verification is superfluous, since we know that the kernel of μ is locally free (see Lemma 8, §4), and thus corresponds to a subbundle N of E'^* such that $E'^*/N = L^*$, and thus necessarily represents the same element as F^* in $K(Y')$. We thus have the exact sequence

$$- \rightarrow \mathcal{O}_{Y'}(F^*) \rightarrow \mathcal{I}'/\mathcal{I}'^2 \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} \xrightarrow{\mu} \mathcal{I}'/\mathcal{I}'^2 \rightarrow 0. \quad (1)$$

By linearity, and Theorem 2, it suffices to prove part (c) of Lemma 19 in the case where $y = \mathcal{G}$ is a locally free sheaf. In this case, $g^1(y) = \mathcal{G} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'}$ is locally free over \mathcal{O}_Y , and $g^1(y) \cdot \lambda_{-1}F^*$ is the alternating sum of the

$$\mathcal{G} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} \otimes_{\mathcal{O}_{Y'}} (\lambda^i F^*) = \mathcal{G} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} (\lambda^i F^*).$$

Also, $f^! i_!(y)$ is the alternating sum of the $\text{Tor}_i^{\mathcal{O}_X}(\mathcal{G}, \mathcal{O}_{X'})$; it thus suffices to prove that

$$\text{Tor}_i^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) = \lambda^i \text{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \quad (\text{for } i \geq 1) \quad (2)$$

$$\mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) = \mathcal{O}_{Y'}(F^*) \quad (3)$$

$$\mathrm{Tor}_j^{\mathcal{O}_X}(\mathcal{G}, \mathcal{O}_{X'}) = \mathcal{G} \otimes_{\mathcal{O}_Y} \mathrm{Tor}_j^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \quad (\text{for } j \geq 1). \quad (4)$$

The equalities (3) and (4) correspond to the particular case of part (c) of [Lemma 19](#) where $y = 1$; their proof will be analogous to that of [Proposition 12 \(§10\)](#). We will first prove (2), as well as the fact that the $\mathrm{Tor}_i^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'})$ are annihilated by \mathcal{J}' , and can thus be identified with sheaves on Y' .

It is clear that the two sides of (2) are zero on an open subset that does not meet Y' . To prove (2) in the neighbourhood of a point $b' \in Y'$, we reuse the notation from [§10.3](#) and [§12.2](#). So we need to calculate the homology of the complex

$$0 \rightarrow \mathfrak{M}_p \otimes_{\mathcal{O}_X} \mathcal{O}_{X'} \rightarrow \dots \rightarrow \mathcal{O}_{X'}$$

which can also be written as

$$0 \rightarrow \mathcal{O}_{X'} \otimes_k \bigwedge^p V \rightarrow \dots \rightarrow \mathcal{O}_{X'}.$$

Suppose that b' is in the open subset U_j' where $t_j \neq 0$. Then it is clear that this complex of sheaves is isomorphic to the complex

$$0 \rightarrow \mathcal{O}_{X'} \otimes_k \bigwedge^p V' \rightarrow \dots \rightarrow \mathcal{O}_{X'}$$

where V' has a basis (e'_i) , and where the differential is characterised by

$$\begin{aligned} \mathfrak{d}(1 \otimes e'_j) &= f_j \otimes 1 \\ \mathfrak{d}(1 \otimes e'_i) &= (f_i - t_i f_j / t_j) \otimes 1 \end{aligned}$$

and by the fact that it extends to a differential of algebras. In this new complex, it is immediate that the cycles of exterior degree s form $\mathcal{O}_{X'} \otimes_k \bigwedge^s (e'_1, \dots, \widehat{e'_j}, \dots, e'_p)$ and that the boundaries of exterior degree s form the ideal

$$f_j \cdot \mathcal{O}_{X'} \otimes_k \bigwedge^s (e'_1, \dots, \widehat{e'_j}, \dots, e'_p).$$

Consequently,

$$\mathrm{Tor}_i^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \cong \mathcal{O}_{Y'} \otimes_k \bigwedge^i (e'_1, \dots, \widehat{e'_j}, \dots, e'_p),$$

which proves that the Tor_i are sheaves on Y' , and, taking the initial remark of [§10.4](#) into account, also gives (2).

Proof. (of (3)). The exact sequence of Tor, applied to

$$0 \rightarrow \mathcal{J} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_Y \rightarrow 0,$$

gives the exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \rightarrow \mathcal{J} \otimes_{\mathcal{O}_X} \mathcal{O}_{X'} \xrightarrow{g} \mathcal{O}_{X'}.$$

It is clear that, if $b = f(b')$, then $\mathcal{J}'_{b'} = \mathcal{J}_b \cdot \mathcal{O}_{b', X'}$, and so the image of g is \mathcal{J}' , and we obtain the exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \rightarrow \mathcal{J} \otimes_{\mathcal{O}_X} \mathcal{O}_{X'} \rightarrow \mathcal{J}' \rightarrow 0. \quad (5)$$

132

Since $\mathrm{Tor}_1(\mathcal{O}_Y, \mathcal{O}_{X'})$ is a sheaf on Y' , we can write

$$\mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{Y'} = \mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'} = \mathrm{Tor}_1(\mathcal{O}_Y, \mathcal{O}_{X'}).$$

Also, \mathcal{I}' can be identified with $\mathcal{O}_{X'}([Y']^{-1})$, and is locally free, and so $\mathrm{Tor}_1^{\mathcal{O}_{X'}}(\mathcal{I}', \mathcal{O}_{Y'}) = 0$, and the exact sequence of Tor, applied to (5), gives the exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{O}_Y, \mathcal{O}_{X'}) \rightarrow \mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{O}_{Y'} \xrightarrow{g} \mathcal{I}' \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{Y'} \rightarrow 0. \quad (6)$$

It is immediate that the canonical image of $\mathcal{I}^2 \otimes_{\mathcal{O}_X} \mathcal{O}_{Y'}$ (resp. $\mathcal{I}'^2 \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{Y'}$) in $\mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{O}_{Y'}$ (resp. $\mathcal{I}' \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{Y'}$) is zero, whence we have canonical isomorphisms

$$\begin{aligned} \mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{O}_{Y'} &= \mathcal{I} / \mathcal{I}^2 \otimes_{\mathcal{O}_X} \mathcal{O}_{Y'} = \mathcal{I} / \mathcal{I}^2 \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} \\ \mathcal{I}' \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{Y'} &= \mathcal{I}' / \mathcal{I}'^2 \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{Y'} = \mathcal{I}' / \mathcal{I}'^2 \end{aligned}$$

which send g to the homomorphism μ from the sequence (1). The equality (2) then follows from (1) and (6).

The equality (4) then follows from an associativity formula of Tor (see [3]). We consider $T(\mathcal{G}, \mathcal{O}_{X'}) = \mathcal{G} \otimes_{\mathcal{O}_Y} \mathcal{O}_Y \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{X'}$ as a functor in \mathcal{G} and $\mathcal{O}_{X'}$. To calculate its left-derived functors $L_i T$, we have two spectral sequences, with respective E_2 pages

$$\begin{aligned} E_2^{ij} &= \mathrm{Tor}_i^{\mathcal{O}_Y}(\mathrm{Tor}_j^{\mathcal{O}_X}(\mathcal{O}_{X'}, \mathcal{O}_Y), \mathcal{G}) \\ E_2'^{ij} &= \mathrm{Tor}_j^{\mathcal{O}_X}(\mathrm{Tor}_i^{\mathcal{O}_Y}(\mathcal{G}, \mathcal{O}_Y), \mathcal{O}_{X'}). \end{aligned}$$

From this, it evidently follows that $E_2^{ij} = E_2'^{ji} = 0$ if $i > 0$, and thus that

$$E_2^{0j} = E_2'^{j0} = L_j T(\mathcal{G}, \mathcal{O}_{X'}) \square$$

since E_2^{0j} and $E_2'^{j0}$ are equal to the right- and left-hand sides (respectively) of (4), this proves the equality. \square

16 Proof of part (d) of Lemma 19

Lemma 20. *Let G be an ample vector bundle of rank $(q + k)$ on Y ($q = \dim Y$; $k \geq 0$). Then G contains a trivial subbundle of rank k .*

Proof. The fact that G is ample implies that, at each point y , the fibre G_y is generated by its sections. So there exists a finite-dimensional k -vector space V of sections, such that the map $r_y: V \rightarrow G_y$, that sends each section to its value at y , is surjective for all $y \in Y$. (If Y is not complete, take a cover by affine open subsets that are complete, and then extract a finite subcover by “quasi-compactness.”) We thus have an exact sequence

$$0 \rightarrow N_y \rightarrow V \rightarrow E_y \rightarrow 0$$

for all $y \in Y$, where $\mathrm{codim}_V N_y = q + k$. The N_y form a vector subbundle N' of the trivial | p. 133

bundle $V \times Y$, by Lemma 8 (§4). The injection of the N_y into V thus defines a morphism $u: N \rightarrow V$. Since $\dim V = q + \dim N_y$, the closure of $\text{Im } u$ is of codimension $\geq k$. Since $\text{Im } u$ is a union of vector subspaces, $\overline{\text{Im } u}$ is an algebraic homogeneous space; consequently, there exists a subspace W of dimension k whose intersection with $\overline{\text{Im } u}$ is $\{0\}$. Then $W \cap N_y = (0)$ for all $y \in Y$, and so W defines the required subbundle. (For this proof, see [1].) \square

Lemma 21. *Let G be a vector bundle of rank $p = q + k$ on Y ($q = \dim Y$). Then $\lambda^s(G - k) = 0$ for all $s \geq q + 1$.*

Proof. Let h be the bundle associated to a hyperplane section of Y . Then (by the Corollary of Proposition 11)

$$(1 - h)^{q+1} = 0$$

and so $h = 1 + u$, with $u^{q+1} = 0$, whence

$$h^n = \sum_{0 \leq i \leq q} \binom{n}{i} u^i.$$

It follows that

$$\lambda_t(Gh^n - k) = \prod_{i=1}^q \lambda_t(Gu^i)^{\binom{n}{i}} \cdot (1 - t)^{-k}$$

and the reader can, from this, deduce that $\lambda^s(Gh^n - k)$ is of the form

$$\lambda^s(Gh^n - k) = \sum_{i=1}^{m_s} B_{s,i} P_{s,i}(n)$$

where $B_{s,i} \in K(Y)$, and where $P_{s,i}(n)$ is a polynomial with rational coefficients that, for all $n > 0$ large enough, takes integer values. We then know (Hilbert) that $P_{s,i}$ is a \mathbb{Z} -linear combination of the polynomials

$$\binom{X}{j} = X(X-1)\dots(X-j+1)/j!$$

and so, finally,

$$\lambda^s(Gh^n - k) = \sum_{i=0}^{n_s} A_{s,i} \binom{n}{i} \quad (\text{for } A_{s,i} \in K(Y)). \quad (1)$$

For $n > n_0$, the bundle Gh^n is ample ([12]), and so (Lemma 20) contains a trivial bundle of rank $k < q$ and $Gh^n - k$ can be identified with a bundle of rank q ; its s th exterior power is then evidently zero for all $s \geq q + 1$. Given (1), it suffices to show that, if the polynomial

$$P(n) = \sum_{i=0}^m A_i \binom{n}{i} \quad (\text{for } A_i \in K(Y))$$

| p. 134

is zero for $n > n_0$, then all the A_i are zero. For this, we proceed by induction on m , and consider the first difference

$$\begin{aligned}\Delta P(n) &= P(n+1) - P(n) \\ &= \sum_{i=1}^m A_i \binom{n}{i-1} \\ &= \sum_{j=0}^{m-1} A_{j+1} \binom{n}{j}.\end{aligned}$$

Since $\Delta P(n) = 0$ for $n > n_0$, the A_i ($1 \leq i \leq m$) are zero by the induction hypothesis, whence clearly $A_0 = 0$ as well. \square

Lemma 22. *Let G and L be vector bundles on Y , with respective ranks p and 1. Then*

- a. $\lambda^p(G-1) = (-1)^p \lambda_{-1}(G)$;
- b. $\lambda_t G(1-L) \equiv 1 \pmod{1-L}$;

and so, if $G_1 \equiv G_2 \pmod{1-L}$ [$G_1, G_2 \in K(Y)$], then $\lambda^i G_1 \equiv \lambda^i G_2 \pmod{1-L}$ for all $i \geq 1$.

Proof. (of part (d) of Lemma 19). By Lemma 22, we have that

$$(-1)^{p-1} \lambda_{-1} F^* = \lambda^{p-1}(F^* - 1).$$

But $E'^*/F^* = L^*$, so $F^* - 1 \equiv E'^* - 2 \pmod{1-L^*}$, whence (Lemma 22)

$$\lambda_{-1} F^* \equiv \lambda^{p-1}(E'^* - 2) \pmod{1-L^*},$$

which can also be written as

$$\lambda_{-1} F^* \equiv g^!(\lambda^{p-1}(E'^* - 2)) \pmod{1-L^*}.$$

It thus suffices to show that $\lambda^{p-1}(E'^* - 2) = 0$ if $p \geq \dim Y + 2$, which follows from Lemma 21. \square | p. 135

17 An application of R–R

The following application (pointed out by Hirzebruch) is about “integration on the fibre” in an algebraic bundle. We place ourselves in the classical case, i.e. $k = \mathbb{C}$.

Proposition 16. *Let (E, B, F, g) be an algebraic bundle, where E , B , and F are projective, irreducible, and non-singular, and let ξ be the tangent bundle along the fibres. Then*

$$g_*(T(\xi)) = \text{To}(F) \cdot 1.$$

[Here $\text{To}(X)$ denotes the Todd genus of X .]

Proof. The tangent bundle to E is an extension of ξ by the fibre induced by the tangent bundle to B , so

$$T(X) = g^*(T(B)) \cdot T(\xi),$$

whence

$$g_*(T(X)) = T(B) \cdot g_*(T(\xi)).$$

We apply R–R to g and the bundle 1 on X , which gives us that

$$g_*(T(X)) = \text{ch } g_!(1) \cdot T(B),$$

and so, given the above,

$$g_*(T(\xi)) = \text{ch } g_!(1).$$

We need to calculate $g_!(1)$. Let U be an affine open subset of B over which the bundle is trivial. By Künneth (see the proof of Lemma 16), we have that

$$H^q(U \times F, \mathcal{O}_X) = \sum_{i+j=q} H^i(U, \mathcal{O}_U) \otimes H^j(F, \mathcal{O}_F).$$

Since U is affine, $H^i(U, \mathcal{O}_U) = 0$ for $i > 0$, and H^0 can be identified with regular functions on U . But, in an algebraic bundle, the structure group G is connected (by hypothesis), and so acts trivially on $H^j(F, \mathcal{O}_F)$, which is the type-(0, j) component of $H^j(F, \mathbb{C})$.⁴ It thus follows that

$$g_!(1) = \mathcal{O}_B \otimes \left(\sum_q (-1)^q H^q(F, \mathcal{O}_F) \right).$$

Then $g_!(1)$ is the alternating sum of trivial bundles, and so $g_!(1)$ has exactly one component, which is in degree 0, and which is the alternating sum of the $\dim H^q(F, \mathcal{O}_F)$, which proves the proposition. \square

| p. 136

Remark. This proposition implies that the multiplicative sequence that defines the Todd class is “strictly multiplicative” for algebraic fibres, in the terminology of Borel-Hirzebruch [2]. In particular, it thus follows that $\text{To}(E) = \text{To}(B) \cdot \text{To}(F)$. In [2], this property, of strict multiplicativity, is proven in the differentiable, almost-complex case, and where also the fibre is some G/T or associated space.

Bibliography

- [1] M. Atiyah. “Vector bundles over an elliptic curve.” *Proc. London Math. Soc.* **7** (1957), 414–452.
- [2] A. Borel, F. Hirzebruch. “Characteristic classes and homogeneous spaces, II.” *Amer. J. Math.* (to appear).
- [3] H. Cartan, S. Eilenberg. *Homological algebra*. Princeton University Press, 1956. *Princeton Math. Series*.

⁴We do not know if the fact that G acts trivially on the $H^j(F, \mathcal{O}_F)$ remains true in characteristic $p > 0$. This is why we had to assume that $k = \mathbb{C}$.

- [4] C. Chevalley. “La notion de correspondance propre en géométrie algébrique.” *Séminaire Bourbaki*. **10** (n.d.), Talk no. 152.
- [5] S.C. Chevalley. *Anneaux de chow et applications*. 1958. **2**.
- [6] W. Chow. “On equivalence classes of cycles in an algebraic variety.” *Ann. Math.* **64** (1956), 450–479.
- [7] A. Grothendieck. “Sur les faisceaux algébriques et les faisceaux analytiques cohérents.” *Séminaire H. Cartan*. **9** (n.d.), Talk no. 2.
- [8] A. Grothendieck. “Sur quelques points d’algèbre homologique.” *Tohoku Math. J.* **9** (1957), 119–221.
- [9] A. Grothendieck. “La théorie des classes de chern.” *Bull. Soc. Math. France*. **86** (1958), 137–154.
- [10] F. Hirzebruch. *Neue topologische methoden in der algebraischen geometrie*. Springer, 1956. *Ergebnisse Der Mathematik*.
- [11] P. Samuel. “Rational equivalence of arbitrary cycles.” *Amer. J. Math.* **78** (1956), 383–400.
- [12] J.-P. Serre. “Faisceaux algébriques cohérents.” *Ann. Math.* **61** (1955), 197–279.
- [13] J.-P. Serre. “Géométrie algébrique et géométrie analytique.” *Ann. Inst. Fourier, Grenoble*. **6** (1955–56), 1–42.
- [14] J.-P. Serre. “Sur la cohomologie des variétés algébriques.” *J. Math. Pures Et Appl.* **36** (1957), 1–16.