

Divisors in algebraic geometry

C.S. Seshadri

1958–59

Translator’s note

This page is a translation into English of the following:

Seshadri, C. S. “Diviseurs en géométrie algébrique.” *Séminaire Claude Chevalley* 4 (1958–59), Talk no. 4. numdam.org/item/SCC_1958-1959__4__A4_0

The translator (Tim Hosgood) takes full responsibility for any errors introduced, and claims no rights to any of the mathematical content herein.

Version: [94f6dce](#)

In the first part of this talk, we will prove a theorem of Serre on complete varieties [6], following the methods of Grothendieck [4]. The second part is dedicated to generalities on divisors. In the literature, we often call the divisors studied here “locally principal” divisors. | p. 4-01

The algebraic spaces considered here are defined over an algebraically closed field K . By “variety,” we mean an irreducible algebraic space. If X is an algebraic space, we denote by $\mathcal{O}(X)$, $\mathcal{R}(X)$, etc. (or simply \mathcal{O} , \mathcal{R} , etc.) the structure sheaf, of regular functions, etc. on X (to define $\mathcal{R}(X)$ we assume that X is a variety). By “coherent sheaf” on X , we mean a coherent sheaf of \mathcal{O} -modules on X .

1 Preliminaries

References: [4–6]

If M is a module over an integral ring A (commutative and with 1), then we say that an element $m \in M$ is a *torsion element* if there exists some non-zero $a \in A$ such that $a \cdot m = 0$. We say that M is a *torsion module* (resp. *torsion-free module*) if every element of M is a torsion element (resp. if $M \neq 0$ and no non-zero element of M is a torsion element). The torsion elements of M form a torsion submodule of M (denoted by $T(M)$); if $M \neq 0$, then $M/T(M)$ is a torsion-free module. If M is a torsion module of finite type over A , then the ideal $\text{ann} M$ of A (the ideal of A given by the elements $a \in A$ such that $aM = 0$) is non-zero.

Let X be an algebraic space and \mathcal{F} a sheaf of \mathcal{O} -modules on X . We define $\text{supp } \mathcal{F}$ to be the set of points $x \in X$ such that $\mathcal{F}_x \neq 0$. If \mathcal{F} is coherent, then $\text{supp } \mathcal{F}$ is a closed subset of X . If X is affine, then $\text{supp } \mathcal{F}$ is the set defined by the ideal $\text{ann} H^0(X, \mathcal{F})$ of the affine algebra $H^0(X, \mathcal{O})$, where $H^0(X, \mathcal{F})$ is considered as a module over $H^0(X, \mathcal{O})$.

A sheaf \mathcal{F} of \mathcal{O} -modules on a variety X is said to be a *torsion sheaf* (resp. *torsion-free sheaf*) if, for every $x \in X$, the module \mathcal{F}_x over the ring \mathcal{O}_x is a torsion module (resp. torsion-free module).

Proposition 1. *If \mathcal{F} is a coherent sheaf on a variety X , then there exists a coherent subsheaf $T(\mathcal{F})$ of \mathcal{F} (and only one) such that $(T(\mathcal{F}))_x = T(\mathcal{F}_x)$.*

Proof. The uniqueness is trivial. The existence is a consequence of the fact that, if X is affine, then $T(\mathcal{F}_x)$ is given by localisation of the module $T(\mathrm{H}^0_*(X, \mathcal{F}))$ with respect to the maximal ideal of $\mathrm{H}^0(X, \mathcal{O})$ that defines x . \square

Corollary *If $\mathcal{F} \neq 0$ then $\mathcal{F}/T(\mathcal{F})$ is a torsion-free coherent sheaf.*¹

Proposition 2. *If \mathcal{F} is a coherent sheaf on the variety X , then $\mathrm{supp} \mathcal{F} \neq X$ if and only if \mathcal{F} is a torsion sheaf.*

Proof. This is a trivial consequence of the fact that, if U is an affine open subset, then $\mathrm{supp} \mathcal{F} \cap U$ is defined by the ideal $\mathrm{ann} \mathrm{H}^0(U, \mathcal{F})$ of $\mathrm{H}^0(U, \mathcal{O})$, where $\mathrm{H}^0(U, \mathcal{F})$ is considered as a module over $\mathrm{H}^0(U, \mathcal{O})$. \square

Proposition 3. *If \mathcal{F} is a torsion-free coherent sheaf on a variety X , with $\mathcal{F} \subset \mathcal{R}^n$, then there exists a coherent sheaf $\mathcal{I} \neq 0$ of ideals of \mathcal{O} such that $\mathcal{I} \cdot \mathcal{F} \subset \mathcal{O}^n$.*

Proof. Let \mathcal{I}_x be the ideal $[\mathcal{O}_x^n : \mathcal{F}_x]$ of \mathcal{O}_x , i.e. the ideal of elements i_x of \mathcal{O}_x such that $i_x \mathcal{F}_x \subset \mathcal{O}_x^n$. Since \mathcal{F}_x is of finite type over \mathcal{O}_x , we know that $\mathcal{I}_x \neq 0$. If we take an affine open subset U of X , then we can prove that \mathcal{I}_x is given by localisation of the ideal $[\mathrm{H}^0(U, \mathcal{O}^n) : \mathrm{H}^0(U, \mathcal{F})]$ of $\mathrm{H}^0(U, \mathcal{O})$ by the maximal ideal of $\mathrm{H}^0(U, \mathcal{O})$ that defines x . Thus $\{\mathcal{I}_x\}_{x \in X}$ defines a coherent sheaf \mathcal{I} of ideals of \mathcal{O} such that $\mathcal{I} \cdot \mathcal{F} \subset \mathcal{O}^n$. \square

Let \mathcal{F} be a torsion-free coherent sheaf on a variety X . Then the canonical homomorphism $\mathcal{F} \rightarrow \mathcal{F} \otimes_{\mathcal{O}} \mathcal{R}$ is injective. The sheaves \mathcal{R} and $\mathcal{F} \otimes_{\mathcal{O}} \mathcal{R}$ are locally constant sheaves, and thus constant ([5]). We can then identify $\mathcal{F} \otimes_{\mathcal{O}} \mathcal{R}$ with a vector space of finite dimension over \mathcal{R} (we identify the field of rational functions with the sheaf \mathcal{R} since \mathcal{R} is constant). We call this dimension the *rank* of \mathcal{F} , and we can then consider \mathcal{F} as a subsheaf of \mathcal{R}^n , where $n = \mathrm{rank} \mathcal{F}$.

Proposition 4. *Under the same hypotheses as in Proposition 3, there exists a coherent sheaf $\mathcal{I} \neq 0$ of ideals of \mathcal{O} such that $\mathcal{I} \cdot \mathcal{F} \subset \mathcal{O}^n$, where $n = \mathrm{rank} \mathcal{F}$; then $\mathcal{O}^n/(\mathcal{I} \cdot \mathcal{F})$ and $\mathcal{F}/(\mathcal{I} \cdot \mathcal{F})$ are torsion sheaves.*

Proof. The proof is immediate. \square

If Y is a closed subset of an algebraic space X , then we denote by \mathcal{I}_Y the coherent sheaf of ideals of \mathcal{O} defined by Y .

Proposition 5. *Let Y be a closed subset of an algebraic space X , and \mathcal{F} a coherent sheaf on X , with $\text{supp } \mathcal{F} \subset Y$; then there exists an integer k such that $\mathcal{I}_Y^k \mathcal{F} = 0$.*

Proof. We can reduce to the case where X is affine, since there exists a finite cover of X by affine opens. In this case, the hypothesis implies that the set defined by the ideal $\text{ann}H^0(X, \mathcal{F})$ is contained in Y . This implies, as is well known, that $\text{ann}H^0(X, \mathcal{F}) \supset \mathcal{I}_Y^k$. \square

Proposition 6. *Let \mathcal{F} be a coherent sheaf of fractional ideals on a variety X (i.e. a coherent subsheaf of \mathcal{R}) such that, for every x outside of a closed subset Y of X , \mathcal{F}_x is an ideal of \mathcal{O}_x . Then there exists an integer k such that $\mathcal{I}_Y^k \cdot \mathcal{F} \subset \mathcal{O}$.*

Proof. By [Proposition 3](#) and the hypothesis, there exists a coherent sheaf \mathcal{J} of ideals of \mathcal{O} such that $\mathcal{J}_x = \mathcal{O}_x$ if $x \notin Y$, and such that $\mathcal{J} \cdot \mathcal{F} \subset \mathcal{O}$. Thus $\text{supp}(\mathcal{O}/\mathcal{J}) \subset Y$, and, by [Proposition 5](#), there exists an integer k such that $\mathcal{I}_Y^k(\mathcal{O}/\mathcal{J}) = 0$. This implies that $\mathcal{I}_Y^k \subset \mathcal{J}$. \square

2 Dévissage theorem

Let \mathcal{C} be an abelian category, and \mathcal{C}' a subcategory of objects of \mathcal{C} . We say that \mathcal{C}' is *left exact in \mathcal{C}* if²

1. every subobject of an object of \mathcal{C}' is in \mathcal{C}' ;
2. for every exact sequence $0 \rightarrow \mathcal{A}' \rightarrow \mathcal{A} \rightarrow \mathcal{A}'' \rightarrow 0$ in \mathcal{C} , the object \mathcal{A} is in \mathcal{C}' if the other two objects are in \mathcal{C}' .

Let X be an algebraic space. We denote by $\mathcal{C}(X)$ the abelian category of coherent sheaves on X . If Y is a closed subset of X , then a coherent sheaf on Y has a canonical extension to a coherent sheaf on X (extending by 0 outside of Y), and so we can consider $\mathcal{C}(Y)$ as a subcategory of $\mathcal{C}(X)$. With this notation, we have the following theorem: | p. 4-04

Theorem (Dévissage). *Let \mathcal{D} be a left-exact subcategory of $\mathcal{C}(X)$ that has the following property: for every closed irreducible subset Y of X , there exists a coherent sheaf \mathcal{M}_Y of $\mathcal{C}(Y)$ that belongs to \mathcal{D} , and that is torsion-free as a sheaf on Y . Then $\mathcal{D} = \mathcal{C}(X)$.*

¹[Trans.] The condition that $\mathcal{F} \neq 0$ is unnecessary, but we include it here since it is in the original. Note that the zero sheaf is indeed a torsion-free sheaf, otherwise any coherent torsion sheaf \mathcal{F} provides a counterexample to this corollary.

²The axioms here that define a left-exact subcategory are slightly stronger than those of Grothendieck [4].

Proof. The proof works by induction on the dimension of X . If $\dim X = 0$, then X consists of a finite number of points P_1, \dots, P_r , and a coherent sheaf on X can be identified with a system $\{N_i\}_{i=1, \dots, r}$, where N_i is a vector space of finite dimension over K . Thus the sheaf \mathcal{M}_{P_i} on P_i that we have, by hypothesis, is a vector space of finite dimension over K . By the axioms of a left-exact subcategory, it is trivial to show that every system $\{N_i\}_{i=1, \dots, r}$, where N_i is a vector space of finite dimension over K , considered as a coherent sheaf on X , belongs to \mathcal{D} .

Now assume that we have proven the theorem for all dimensions $\leq (n-1)$. Let $\dim X = n$. Let Y be a closed subset of X such that $\dim Y \leq (n-1)$. We can easily show that $\mathcal{D} \cap \mathcal{C}(Y)$ is a left-exact subcategory of $\mathcal{C}(Y)$ that satisfies the hypotheses of the theorem. So, by the induction hypothesis, $\mathcal{D} \supset \mathcal{C}(Y)$.

We will now prove that, if \mathcal{F} is a coherent sheaf on X with $\text{supp } \mathcal{F} = Y$, then $\mathcal{F} \in \mathcal{D}$. If $\mathcal{I}_Y \cdot \mathcal{F} = 0$, then $\mathcal{F} \in \mathcal{C}(Y)$, and, by the above, $\mathcal{F} \in \mathcal{D}$. No matter what, by [Proposition 5](#), there exists an integer $k \geq 1$ such that $\mathcal{I}_Y^k \mathcal{F} = 0$. We will complete the proof by induction on k . Suppose that that claim has been proven for every coherent sheaf \mathcal{G} on X such that $\mathcal{I}_Y^{k-1} \mathcal{G} = 0$. For \mathcal{F} , we have an exact sequence

$$0 \rightarrow \mathcal{I}_Y \cdot \mathcal{F} \rightarrow \mathcal{F} \rightarrow \mathcal{F}/(\mathcal{I}_Y \cdot \mathcal{F}) \rightarrow 0.$$

The sheaf $\mathcal{I}_Y \mathcal{F}$ is annihilated by \mathcal{I}_Y^{k-1} , and the sheaf $\mathcal{F}/(\mathcal{I}_Y \mathcal{F})$ is annihilated by \mathcal{I}_Y . Thus $\mathcal{I}_Y \mathcal{F}$ and $\mathcal{F}/(\mathcal{I}_Y \mathcal{F})$ belong to \mathcal{D} . This implies that $\mathcal{F} \in \mathcal{D}$.

Suppose that X is a variety, and that \mathcal{F} is a torsion-free sheaf on X . We can consider \mathcal{F} as a coherent subsheaf of \mathcal{R}^n , where $n = \text{rank } \mathcal{F}$, and, by [Proposition 4](#), there then exists a coherent sheaf of ideals \mathcal{I} such that $\mathcal{I} \cdot \mathcal{F} \subset \mathcal{O}^n$, and such that the sheaves $\mathcal{F}/(\mathcal{I} \mathcal{F})$ and $\mathcal{O}^n/(\mathcal{I} \mathcal{F})$ are torsion sheaves. Since $\mathcal{F}/(\mathcal{I} \mathcal{F})$ is a torsion sheaf, $\mathcal{F}/(\mathcal{I} \mathcal{F}) \in \mathcal{D}$; thus $\mathcal{F} \in \mathcal{D}$ if and only if $\mathcal{I} \mathcal{F} \in \mathcal{D}$. Analogously, $\mathcal{I} \mathcal{F} \in \mathcal{D}$ if and only if $\mathcal{O}^n \in \mathcal{D}$, and, by the axioms of an exact subcategory, if and only if $\mathcal{O} \in \mathcal{D}$. Thus $\mathcal{F} \in \mathcal{D}$ if and only if $\mathcal{O} \in \mathcal{D}$. If we repeat the same argument for the torsion-free sheaf \mathcal{M}_X , which we have by hypothesis, then we see that $\mathcal{O} \in \mathcal{D}$, which implies that $\mathcal{F} \in \mathcal{D}$. | p. 4-05

Suppose again that X is a variety, but now that \mathcal{F} is an arbitrary coherent sheaf. We will show that $\mathcal{F} \in \mathcal{D}$. We can assume that $\mathcal{F} \neq 0$, and we then have

$$0 \rightarrow T(\mathcal{F}) \rightarrow \mathcal{F} \rightarrow \mathcal{F}/T(\mathcal{F}) \rightarrow 0$$

where $T(\mathcal{F})$ is a torsion sheaf, and $\mathcal{F}/T(\mathcal{F})$ is a torsion-free sheaf. By [Proposition 2](#), $\text{supp } T(\mathcal{F}) \neq X$, and, since X is a variety, $\dim \text{supp } T(\mathcal{F}) < \dim T(X)$. We then have, by the induction hypothesis, that $T(\mathcal{F}) \in \mathcal{D}$, and we have just proven that $\mathcal{F}/T(\mathcal{F}) \in \mathcal{D}$. Thus $\mathcal{F} \in \mathcal{D}$.

Now let X be an arbitrary algebraic space, and X_1, \dots, X_p its irreducible components. If \mathcal{F} is a coherent sheaf on X , then $\mathcal{F}/(\mathcal{I}_{X_i} \mathcal{F})$ can be identified with a sheaf on the variety X_i (where \mathcal{I}_{X_i} is the sheaf of ideals of $\mathcal{O}(X)$ determined by X_i), and, by the above, $\mathcal{F}/(\mathcal{I}_{X_i} \mathcal{F}) \in \mathcal{D}$. Thus the sheaf $\mathcal{G} = \sum_{i=1}^p \mathcal{F}/(\mathcal{I}_{X_i} \mathcal{F})$ belongs to \mathcal{D} . We have a canonical homomorphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$. The image of φ is a coherent subsheaf of \mathcal{G} , and so the image of φ belongs to \mathcal{D} .

We know that $\text{supp } \text{Ker } \varphi \subset \bigcup_{i \neq j} X_i \cap X_j$, and so $\dim \text{supp } \text{Ker } \varphi < \dim X$, and, by the induction hypothesis, $\text{Ker } \varphi \in \mathcal{D}$. Thus $\mathcal{F} \in \mathcal{D}$, and the theorem is proven. \square

Corollary (Serre's Theorem). *If \mathcal{F} is a coherent sheaf on a complete algebraic space X , then $H^0(X, \mathcal{F})$ is a vector space of finite dimension over K .*

Proof. We take \mathcal{D} to be the category of all coherent sheaves \mathcal{F} on X such that $H^0(X, \mathcal{F})$ is of finite dimension over K . We can prove that \mathcal{D} is a left-exact subcategory of $\mathcal{C}(X)$. Also, we know that, if Y is an irreducible closed subset of X , then Y is a complete variety. Thus the coherent sheaf $\mathcal{O}(Y)$ on Y is a torsion-free sheaf with the property that $H^0(Y, \mathcal{O}(Y)) \cong K$, and so $H^0(X, \mathcal{O}(Y)) = H^0(Y, \mathcal{O}(Y))$ is of finite dimension over K (we denote also by $\mathcal{O}(Y)$ the canonical extension of $\mathcal{O}(Y)$ to X). By the Theorem, the corollary is proven. \square

3 Divisors (Generalities)

Let X be an algebraic variety, and $\mathcal{R}^\times(X)$ and $\mathcal{O}^\times(X)$ (or simply \mathcal{R}^\times and \mathcal{O}^\times) the constant sheaf on X of non-zero rational functions and the sheaf on X of invertible regular functions (respectively). The sheaves \mathcal{R}^\times and \mathcal{O}^\times , endowed with their multiplicative structure, are sheaves of abelian groups. | p. 4-06

A *divisor* D on X is a section of the quotient sheaf $\mathcal{R}^\times/\mathcal{O}^\times$. An element of \mathcal{R}^\times that is a representative of the value $D(x)$ of D at x is called a *definition function of D at x* . More generally, a function $f \in \mathcal{R}^\times$ is called a *definition function of D in an open subset U* if, for all $x \in U$, f is a representative of $D(x)$; then f is determined up to an invertible regular function on U . Since we can locally lift a section of $\mathcal{R}^\times/\mathcal{O}^\times$ to a section of \mathcal{R}^\times , a divisor D is determined by the following data: a cover $\{U_i\}$ by open subsets, and non-zero rational functions f_i on U_i such that, on $U_i \cap U_j$, $f_{ij} = f_i/f_j$ is an invertible regular function. We have that $f_{ij}f_{jk}f_{ki} = 1$ on $U_i \cap U_j \cap U_k$, and, as is well known, this allows us to construct a locally trivial fibre bundle with K^\times as the structure group; it is easy to see that this fibre bundle is determined up to equivalence [7]. We also know that the coherent sheaves of fractional ideals (i.e. the coherent subsheaves of \mathcal{R}) that are generated by f_i and f_j (respectively) agree on $U_i \cap U_j$, and do not depend on the choice of definition functions of D in U_i and U_j . This implies that the divisor D canonically determines a coherent sheaf of locally principal fractional ideals. We can easily see that the converse is true, and this gives us an equivalent definition of a divisor [1].

A divisor D is said to be *positive* if, for each $x \in X$, $D(x) \in \mathcal{O}_x/\mathcal{O}_x^\times$ (i.e. if all the definition functions of D at x are regular functions in x).

Since $\mathcal{R}^\times/\mathcal{O}^\times$ is a sheaf of abelian groups on X , there is a canonical structure of an abelian group on the set of divisors on X ; this group is called the *group of divisors on X* . The composition law in this group is written additively, and the identity element in this group is thus called the *zero divisor*, and is denoted by (0) .

If f is a non-zero rational function on X , then it defines a divisor $\text{div} f$ by the data $(\text{div} f)(x) = \text{Im} f \subseteq \mathcal{R}^\times/\mathcal{O}_x^\times$. The divisors obtained in this way are called *principal divisors*, and form a subgroup of the group of divisors on X ; the quotient group is called the *group of classes of divisors on X* . Two divisors D_1 and D_2 are said to be equivalent if they are equivalent module the group of principal divisors; we write $D_1 \sim D_2$. We have seen that a divisor defines, up to equivalence, a locally trivial algebraic fibre bundle with structure group K^\times . On the other hand, it is easy to see that a locally trivial algebraic fibre bundle with K^\times as its structure group defines, up to equivalence, a divisor [7]. Thus the group of | p. 4-07

classes of divisors on X is equal to $H^1(X, \mathcal{O}^\times)$, the group of classes of equivalent algebraic fibre bundles with K^\times as their structure group.

We can define, in an analogous way, an *additive divisor* on a variety X as a section of the sheaf \mathcal{R}/\mathcal{O} (the divisors defined above are called *multiplicative divisors*, or simply *divisors*). The additive divisors form an abelian group, and even a vector space over K . An additive divisor is determined by the following data: a cover $\{U_i\}$ of X by open subsets, and rational functions f_i on U_i such that $f_{ij} = f_i - f_j$ is a regular function on $U_i \cap U_j$. We can define, as for (multiplicative) divisors, the notions of definition functions of an additive divisor, equivalence between two additive divisors, etc. We find, for example, that $H^1(X, \mathcal{O})$ is equal to the group of classes of additive divisors on X .

Let D be a (multiplicative) divisor on X . We define $\text{supp} D$ to be the set of points $x \in X$ such that $D(x)$ is not the identity element in $\mathcal{R}^\times/\mathcal{O}_x^\times$, i.e. such that every definition function of D at x is either not defined at x or takes the value 0 at x .

Proposition 7. *The support of a divisor D on a variety X is a closed subset $\neq X$ of X , and $D = 0$ if and only if the support is empty.*

Proof. The latter claim is trivial. For the former, we prove that the set E of points $x \in X$ such that every definition function of D at x belongs to \mathcal{O}_x^\times is a non-empty open subset; indeed, if we take a definition function g of D at x , then it is also a definition function of D in an open subset U that contains x . By hypothesis, if $x \in E$, then g is regular at x and $g(x) \neq 0$, and we can choose U such that g is regular and invertible on U , which proves that E is open. \square

| p. 4-08

Proposition 8. *If D is a divisor on a normal variety X , then $\text{supp} D$ is a union of hypersurfaces (i.e. of closed subvarieties of codimension 1).*

Proof. If f is a function on a normal variety Y , we know that, if f is not defined at $x \in Y$, then x belongs to a variety of poles or zeros of f (i.e. to an irreducible component of the closure of the set of points $x \in Y$ such that $f(x) \in \{0, \infty\}$). So, if we take f to be a definition function for D in some open subset $U \subset X$, then $\text{supp} D \cap U$ is the union of the pole and zero varieties of f in U , and we know that these varieties are of codimension 1 ([2, chapitre III]). \square

Remark. If X is not normal, then the support of a divisor D on X is not necessarily of codimension 1. It is easy to define an affine variety X of dimension > 1 that is normal everywhere except at a single point x_0 (for example, the point (a, ab, b^2, b^3) in the four-dimensional space K^4). There exists a function u that is everywhere defined on X , and which is entire on the local ring of x_0 , but which is not contained in this ring; by adding, if necessary, a constant, we can assume that x_0 is not a zero of u . There is then an open neighbourhood X' of x_0 such that the divisor of the function induced by u on X' has support equal to the single point x_0 .

Suppose that X is a normal variety, and D is a divisor on X . Let S be a hypersurface of X . If f is a definition function of D at $x \in S$, then the order of f on S ([2]) does not depend on the choice of f , nor on $x \in S$. We denote this integer by $\text{ord}_S D$. It is easy

to see that $\text{ord}_S D = 0$ if and only if $S \not\subset \text{supp} D$. If we now take the formal combination $C = \sum_S (\text{ord}_S D) \cdot S$, where S runs over the set of all hypersurfaces of X , then C is a cycle of codimension 1, and we call it the *associated cycle* of the divisor D .

Proposition 9. *Let X be a normal variety. The map that sends a divisor D to its associated cycle of codimension 1 is an injective homomorphism from the group of divisors on X to the group of cycles of codimension 1.*

Proof. The proof is trivial. □

Proposition 10. *If X is further a non-singular variety, then the homomorphism that sends a divisor to its associated cycle is bijective.*

| p. 4-09

Proof. It suffices to show that, for every hypersurface S , there exists a divisor D such that the cycle $1 \cdot S$ is the cycle associated to D . Since X is non-singular, for every $x \in X$, the local ring \mathcal{O}_x is factorial [3]; thus, for every $x \in S$, S is defined by one single equation in a neighbourhood of x . So there exists a cover $\{U_i\}_{i=1, \dots, p}$ of S by open subsets U_i of X , and, for each i , a regular function f_i on U_i that is non-zero outside of $U_i \cap S$ in U_i with $\text{ord}_S f_i = 1$. It then follows that f_i/f_j is an invertible regular function in $U_i \cap U_j$. Now take the cover $\{U_i\}_{i=0, \dots, p}$, where $U_0 = CS$, and take $f_0 = 1$. It is easy to see that the divisor D for which f_i is a definition function of D on U_i is such that the cycle associated to D is $1 \cdot S$. So the proposition is proven. □

Remark. Proposition 10 is not necessarily true if X is non-singular. For example, for the cone $xy - zw = 0$ in K^4 , the cycle defined by $x = z = 0$ is not a cycle associated to any divisor.

Bibliography

- [1] P. Cartier. “Diviseurs et dérivations en géométrie algébrique.” *Bull. Soc. Math. France.* (1958).
- [2] C. Chevalley. *Fondements de la Géométrie algébrique.* Paris, Secrétariat mathématique, 1958.
- [3] R. Godement. *Propriétés analytiques des localités.* 1955-56. **8**.
- [4] A. Grothendieck. *Sur les faisceaux algébriques et les faisceaux analytiques cohérents.* 1956-57. **9**.
- [5] J.-P. Serre. “Faisceaux algébriques cohérents.” *Ann. Math.* **61** (1955), 197–279.
- [6] J.-P. Serre. “Sur la cohomologie des variétés algébriques.” *J. Math. Pures Et Appl.* **36** (1957), 1–16.
- [7] A. Weil. *Fibre spaces in algebraic geometry.* University of Chicago, 1955.