
On formal complex spaces

Jürgen Bingener

1978

Translator's note

This page is a translation into English of the following:

Bingener, J. "Über formale komplexe Räume." *manuscripta mathematica* **24** (1978), 253–293. DOI: [10.1007/BF01167833](https://doi.org/10.1007/BF01167833).

The translator (Tim Hosgood) takes full responsibility for any errors introduced, and claims no rights to any of the mathematical content herein.

Version: [94f6dce](#)

| p. 253

Abstract. Formal complex spaces, introduced by Krasnov [24] and independently by the author, are the analytic analogues of the formal schemes of Zariski and Grothendieck. Special cases are the formal completions of complex spaces along analytic sets, see Banica [3]. The technique of formal complex spaces has proved to be a useful tool in analytic geometry and allows even applications to purely algebraic problems, see [24], [4] and [7]. Here the basic theory of these spaces is developed: coherence of the structure sheaf, description of the coherent modules, Grauert's coherence theorem for proper maps. . . . We further study the question of exactness of the formal Dolbeault and de Rham complexes.

Introduction

In 1958, Grothendieck introduced formal schemes in algebraic geometry, following on from earlier ideas by Zariski. Since then, the theory of formal schemes has become an important tool in algebraic geometry, cf. e.g. [2,13,20]. Formal structures appeared in (global) analytic geometry for the first time in Grauert's comparison theorem; this is clearly expressed in the proof given in [3]. In [24], Krasnov then explicitly introduced formal complex spaces, and, in particular, formal complex manifolds, and used this to prove theorems about modifications of complex manifolds.

| p. 254

In the present article, we first develop the basic theory of formal complex spaces. These are introduced in §1 as inductive limits of a suitable system of complex spaces. We obtain special formal complex spaces if we consider the formal completions of complex spaces along analytical subsets. Of course, every complex space is also a formal complex space. The structure sheaf \mathcal{O}_X of a formal complex space X is always coherent with local Noetherian stalks ((1.1) and (1.4)). It is important to know that, as in the case of complex spaces,

the **!!TO-DO: intersection rings?!!** of compact Stein subsets of X are excellent Noetherian rings ((1.4) and (1.10)). Formal complex spaces can be (locally) embedded into **!!TO-DO: number spaces?!!**, cf. (1.7).

Formal complex manifolds, i.e. formal complex spaces whose stalks are all regular, are studied in §2. Every point of a formal manifold has an open neighbourhood that is isomorphic to the formal completion of an open subspace of \mathbb{C}^n along an analytic subset. A formal Dolbeault complex can be defined for any formal complex space, and, in the case of formal manifolds, this is a fine resolution of the structure sheaf. This follows directly from the following statement, which is the main result of §2:

Lemma. *Let Y be an open subset of \mathbb{C}^n , $S \subseteq Y$ an analytic set in Y , and \widehat{Y} the formal completion of Y along S . Let $\mathcal{E}_Y^{0,\bullet}$ be the ordinary Dolbeault complex of Y , $\mathcal{I}^{(\infty)}(S)$ the ideal of functions of \mathcal{E}_Y that are flat on S , and $\mathcal{E}_{\widehat{Y}}^{0,\bullet} := \mathcal{E}_Y^{0,\bullet} / \mathcal{I}^{(\infty)}(S) \mathcal{E}_Y^{0,\bullet}$ the formal Dolbeault complex of Y . Then the sequence*

$$0 \rightarrow \mathcal{O}_{\widehat{Y}} \rightarrow \mathcal{E}_{\widehat{Y}} \xrightarrow{\bar{\partial}} \mathcal{E}_{\widehat{Y}}^{0,1} \xrightarrow{\bar{\partial}} \dots \rightarrow \mathcal{E}_{\widehat{Y}}^{0,n} \rightarrow 0$$

is exact.

The fact that $\mathcal{O}_{\widehat{Y}} = \text{Ker}(\mathcal{E}_{\widehat{Y}} \xrightarrow{\bar{\partial}} \mathcal{E}_{\widehat{Y}}^{0,1})$ is exact has already been shown by Krasnov. To prove this, we reduce the problem, using a resolution of singularities and Hironaka’s vanishing theorem, to the case where S is a normal crossing divisor. In this special case, the problem can be solved by “concrete” calculation using theorems of Malgrange.

| p. 255

An analogous statement can be made for the formal de Rham complex. In fact, the following more general statement (cf. (2.11)) holds:

Lemma. *Let S be an analytic set in a complex space Y , defined by a coherent sheaf of \mathcal{O}_Y -ideals \mathcal{I} , and let $\Omega_{\widehat{Y}}^{\bullet} := \varinjlim_k \Omega_Y^{\bullet} / \mathcal{I}^{k+1} \Omega_Y^{\bullet}$. If $Y \setminus S$ is non-singular, then, for all $n \in \mathbb{N}$, the canonical homomorphism*

$$\mathcal{H}^n(\Omega_Y^{\bullet} | S) \rightarrow \mathcal{H}^n(\Omega_{\widehat{Y}}^{\bullet})$$

is bijective.

In the special case where Y is a manifold, then $\Omega_{\widehat{Y}}^{\bullet}$ is a resolution of the constant sheaf \mathbb{C}_S . This latter claim can also be found in Hartshorne [16].

In §3, proper maps between formal complex spaces are considered. For such maps, Grauert’s coherence law (cf. (3.1)) applies. In §4 we show how the most important statements of the relative comparison theory between algebraic and analytic geometry [5,15] can be transferred to the case where the base is a formal complex space.

In §5 we use the results of §4 to study formal meromorphic functions. We mention here the following statement (cf. (5.2)):

Corollary. *Let T be a connected exceptional analytic set in a normal complex space X , let $X \rightarrow Y$ the associated contraction of T to a point $y \in Y$, and let \widehat{X} be the formal completion of X along T . Then the ring $M(\widehat{X})$ of meromorphic functions on \widehat{X} agrees with the quotient field $\mathbb{Q}(\widehat{\mathcal{O}}_{Y,y})$ of the completion of the stalks $\mathcal{O}_{Y,y}$ of y in Y .*

In addition, we determine the ring of meromorphic functions on the product of a normal formal complex space Y with a compact complex algebraic space Z (cf. (5.3)). In the case where Z is the complex projective space $\mathbb{P}_{\mathbb{C}}^r$, we obtain a corollary of Andreotti–Stoll [1, Theorem 6.9].

| p. 256

Some results from the present article have already been used in [4] and [7].

1 Formal complex spaces

Let $X = (X, \mathcal{O}_X)$ be a locally ringed space. For $x \in X$, let \mathfrak{m}_x be the maximal ideal of the stalk \mathcal{O}_x of X at the point x . We denote by \mathcal{I}_X the \mathcal{O}_X ideal whose sections over an open subset U of X are exactly the elements of $\Gamma(U, \mathcal{O}_X)$ such that $f_x \in \mathfrak{m}_x$ for all $x \in U$.

Definition.

A locally ringed space $X = (X, \mathcal{O}_X)$ over $\text{Spec} \mathbb{C}$ is called a *formal complex space* if the following conditions are satisfied:

1. $X_n := (X, \mathcal{O}_X / \mathcal{I}_X^{n+1})$ is a complex space for all $n \in \mathbb{N}$; and
2. the canonical homomorphism $\mathcal{O}_X \rightarrow \varprojlim_n \mathcal{O}_X / \mathcal{I}_X^{n+1}$ is bijective.

Formal complex spaces form a category, with \mathbb{C} -morphisms as the morphisms. If $f : X \rightarrow Y$ is a morphism between formal complex spaces, then the associated homomorphisms $\mathcal{O}_{Y, f(x)} \rightarrow \mathcal{O}_{X, x}$ are local. In particular, we have that $\mathcal{I}_Y \mathcal{O}_X \subseteq \mathcal{I}_X$. If X is a complex space, then $\mathcal{I}_X = \mathcal{N}_X$ is the sheaf of nilpotent elements of X , and so every complex space is also a formal complex space.

As usual, we say that a subset of a complex space is Stein compact if it is a compact, semi-analytic subset that has an neighbourhood system of open Stein sets. This definition can be extended in a trivial way to formal complex spaces: A subset K of a formal complex space X is said to be *Stein compact* if K , regarded as a subset of the complex space X_n , is Stein compact, for all n . Because a complex space is Stein if and only if its reduction is, this condition only needs to be checked for $n = 0$.

The following lemma is fundamental for what follows.

Lemma 1.1.

Let $X = (X, \mathcal{O}_X)$ be a formal complex space, and $K \subseteq X$ a Stein compact subset. Then the following hold:

1. $B_K := \varprojlim_n \Gamma(K, \mathcal{O}_{X_n})$ is a Noetherian ring, which is further separated and complete with respect to the topology defined by the ideal $\mathfrak{b}_K := \text{Ker}(B_K \rightarrow \Gamma(K, \mathcal{O}_{X_0}))$.
2. If $L \subseteq X$ is another Stein compact subset such that $K \subseteq L$, then the canonical homomorphism $B_L \rightarrow B_K$ is flat.
3. \mathcal{O}_X is a coherent sheaf of rings.

Proof. We write $B_{K,n} = \Gamma(K, \mathcal{O}_{X_n})$ and $\mathfrak{b}_n := \Gamma(K, \mathcal{I}_X^n / \mathcal{I}_X^{n+1})$. □

Bibliography

- [1] A. Andreotti, W. Stoll. *Analytic and Algebraic Dependence of Meromorphic Functions*. Springer, 1971. *Lec. Notes Math.* **234**.
- [2] M. Artin. "Algebraization of formal moduli: II. Existence of modifications." *Ann. Math.* **91** (1970), 88–135.
- [3] C. Banica. "Le complété formel d'un espace analytique le long d'un sous-espace: Un théorème de comparaison." *Manuscripta Math.* **6** (1972), 207–244.

- [4] J. Bingener. “Divisorenklassengruppen der Komplettierungen analytischer Algebren.” *Math. Ann.* **217** (1975), 113–120.
- [5] J. Bingener. “Schemata über steinschen algebren.” *Schriftenreihe Des Mathematischen Instituts Der Universität Münster.* **10** (1976).
- [6] J. Bingener. “Holomorph-prävollständige Resträume zu analytischen Mengen in Stein-schen Räumen.” *J.f.d.r.u.a.M.* **285** (1976).
- [7] J. Bingener. “Über die Divisorenklassengruppen lokaler Ringe.” *Math. Ann.* (1977), 173–179.
- [8] N. Bourbaki. *Algèbre commutative.* Hermann, 1961–67.
- [9] E. Brieskorn. “Die Monodromie der isolierten Singularitäten von Hyperflächen.” *Manuscripta Math.* **2** (1970), 103–161.
- [10] H. Cartan. *Séminaire 1960/61.* 1960–61.
- [11] R. Godement. *Théorie des faisceaux.* Hermann, 1964.
- [12] A. Grothendieck. “Géométrie formelle et géométrie algébrique.” *Séminaire Bourbaki.* **11** (1958–59).
- [13] A. Grothendieck. *Cohomologie locale des faisceaux cohérents et Théorèmes de Lefschetz locaux et globaux (SGA 2).* North-Holland Publishing Company, 1968.
- [14] A. Grothendieck, J. Dieudonné. *Éléments de géométrie algébrique.* Pub. Math. I.H.E.S., 1960–1967. **4,8,11,17,20,24,28,32.**
- [15] M. Hakim. *Topos Annelés et Schémas Relatifs.* Springer, 1972.
- [16] R. Hartshorne. “On the de Rham cohomology of algebraic varieties.” **45** (1975), 5–99.
- [17] M. Herrera, D. Liebermann. “Duality and the De Rham Cohomology of Infinitesimal Neighborhoods.” *Inventiones Math.* **13** (1971), 97–124.
- [18] H. Hironaka. “Resolution of singularities of an algebraic variety over a field of character-istic zero: I,II.” *Ann. Math.* **79** (1964), 109–326.
- [19] H. Hironaka. “Flattening theorem in complex-analytic geometry.” *American J. Of Math.* **97** (1975), 503–547.
- [20] H. Hironaka, H. Matsumura. “Formal functions and formal embeddings.” *J. Math. Soc. Japan.* **20** (1968), 52–82.
- [21] H. Hironaka, H. Rossi. “On the Equivalence of Embeddings of Exceptional Complex Spaces.” *Math. Ann.* **156** (1964), 313–333.
- [22] L. Kaup. “Eine Künnethformel für Fréchetgarben.” *Math. Z.* **97** (1967), 158–168.
- [23] R. Kiehl, J.-L. Verdier. “Ein einfacher Beweis des Kohärenzsatzes von Grauert.” *Math. Ann.* **195** (1971), 24–50.

- [24] B.A. Krasnov. “Formal Modifications. Existence Theorems for modifications of complex manifolds.” *Math. USSR Izvestija*. **7** (1973), 847–881.
- [25] R. Narasimhan. “On the Homology Groups of Stein Spaces.” *Inventiones Math.* **2** (1967), 377–385.
- [26] H.-J. Reiffen. “Das Lemma von Poincaré für holomorphe Differentialformen auf komplexen Räumen.” *Math. Z.* **101** (1967), 269–284.
- [27] G. Scheja, U. Storch. “Differentielle Eigenschaften der Lokalisierungen analytischer Algebren.” *Math. Ann.* **197** (1972), 137–170.
- [28] J.-C. Tougeron. *Idéaux de fonctions différentiables*. Springer, 1972.
- [29] K.-W. Wiegmann. “Einbettungen komplexer Räume in Zahlenräume.” *Inventiones Math.* **1** (1966), 229–242.